

Accepted Manuscript

Entanglement of Hard-Core Bose Gas in Degenerate Levels under Local Noise

Z. Gedik, G. Karpat


PII: S0921-4534(09)00869-7
DOI: [10.1016/j.physc.2009.11.152](https://doi.org/10.1016/j.physc.2009.11.152)
Reference: PHYSC 1250908

To appear in: *Physica C*

Received Date: 10 August 2009
Accepted Date: 23 November 2009

Please cite this article as: Z. Gedik, G. Karpat, Entanglement of Hard-Core Bose Gas in Degenerate Levels under Local Noise, *Physica C* (2009), doi: [10.1016/j.physc.2009.11.152](https://doi.org/10.1016/j.physc.2009.11.152)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Entanglement of Hard-Core Bose Gas in Degenerate Levels under Local Noise

Z. Gedik, G. Karpat

Faculty of Engineering and Natural Sciences, Sabanci University, Tuzla, Istanbul 34956, Turkey

Abstract

Quantum entanglement properties of the pseudo-spin representation of the BCS model is investigated. In case of degenerate energy levels, where wave functions take a particularly simple form, spontaneous breaking of exchange symmetry under local noise is studied. Even if the Hamiltonian has the same symmetry, it is shown that there is a non-zero probability to end up with a non-symmetric final state. For small systems, total probability for symmetry breaking is found to be inversely proportional to the system size.

Key words: Quantum decoherence, spontaneous symmetry breaking, mesoscopic superconductivity

PACS: 03.65.Yz, 03.65.Ta, 74.78.Na

Fermions undergoing BCS type pairing interaction can be treated as hard-core bosons in real or momentum space depending on whether they interact strongly or weakly. In both cases, lattice points (or energy levels) are occupied by pairs or else they are empty. Therefore, the system can be described by pseudo-spin variables [1]. Quantum entanglement and superconducting order parameter of such systems have been found to be closely related [2]. In case of degenerate energy levels, wave functions take a particularly simple form and hence all possible eigenstates can be written down easily [3]. Such a degeneracy can for example be due to parabolic energy bands in mesoscopic and nanoscopic superconducting particles [4].

Recently, it has been shown that exchange symmetry of some entangled states can be spontaneously broken under local noise even if the Hamiltonian has the same symmetry [5]. Considering the symmetry of the initial state and the Hamiltonian, this is a very interesting result. Since BCS model of superconductivity can be written in terms of pseudo-spins or (in the language of quantum information theory) in terms of qubits, one might ask consequences of this kind of symmetry breaking on superconducting state.

Our starting point is a single, d -fold degenerate energy level with N spin 1/2 fermions. Assuming that fermions in the same state are paired, the model Hamiltonian becomes

$$H_0 = -g \sum_{n,n'=1}^d a_{n'\uparrow}^\dagger a_{n'\downarrow}^\dagger a_{n\downarrow} a_{n\uparrow}, \quad (1)$$

where $a_{n\sigma}^\dagger$ ($a_{n\sigma}$) creates (annihilates) a fermion in state n with spin σ . Introducing pseudo-spins \vec{s}_n defined by $s_{n+} = a_{n\uparrow}^\dagger a_{n\downarrow}^\dagger$ and $s_{nz} = (1/2)(a_{n\uparrow}^\dagger a_{n\uparrow} + a_{n\downarrow}^\dagger a_{n\downarrow} - 1)$, we can rewrite the Hamiltonian as

$$H_0 = -gS_+S_- \quad (2)$$

where $\vec{S} = \sum_n \vec{s}_n$ is the total spin and S_\pm are the corresponding

raising and lowering operators. Energy eigenvalues are given by

$$E(s) = -(g/4)(N-s)(2d-s-N+2) \quad (3)$$

where seniority number $s = d - 2S$ can take values $s = 0, 2, 4, \dots, N$ when the number of fermions N is even. The ground state, which is invariant under Cooper pair exchange, has the maximum total- S value $d/2$. Degeneracy of each state can be found easily as [6]

$$\Omega(s) = \sum_{i=0}^{s/2} \left[\binom{d-s+2i}{i} - \binom{d-s+2i}{i-1} \right] \binom{d}{s-2i} 2^{s-2i}. \quad (4)$$

Now, let us consider an exchange symmetric local external noise described by, for $d = 2$,

$$H_1(t) = w_1(t)(s_{1z} \otimes I) + w_2(t)(I \otimes s_{2z}), \quad (5)$$

where $w_1(t)$ and $w_2(t)$ are stochastic noise fields that lead to statistically independent Markov processes satisfying

$$\langle w_n(t) \rangle = 0 \quad (6)$$

$$\langle w_n(t)w_n(t') \rangle = a_n \delta(t-t'). \quad (7)$$

In case of real space pairing, such a noise might originate from an external disturbance localized in space in a region of Cooper pair size. Generalization to arbitrary d is obvious. Let the system be in state $\rho(t=0)$, say the ground state. The time evolution of the system's density matrix can be obtained as

$$\rho(t) = \langle U(t)\rho(0)U^\dagger(t) \rangle, \quad (8)$$

where ensemble averages are evaluated over the two noise fields $w_1(t)$ and $w_2(t)$ and the time evolution operator, $U(t)$, is given by

$$U(t) = \exp[-i \int_0^t dt' H(t')]. \quad (9)$$

In Ref. [5], using Kraus representation [7, 8, 9]

$$\rho(t) = \sum_{\mu=1}^M K_{\mu}(t)\rho(0)K_{\mu}^{\dagger}(t), \quad (10)$$

general time evolution has been evaluated and it has been shown that not all possible final states

$$\frac{K_{\mu}(t)\rho(0)K_{\mu}^{\dagger}(t)}{\text{tr}(K_{\mu}(t)\rho(0)K_{\mu}^{\dagger}(t))} \quad (11)$$

have exchange symmetry. The same result has been reproduced for quantum noise using the decoherence Hamiltonian [10]

$$H_1 = s_{1z} \sum_{k=1}^{N_1} \hbar\omega_{1k}\sigma_{1kz} + s_{2z} \sum_{k=1}^{N_2} \hbar\omega_{2k}\sigma_{2kz}. \quad (12)$$

where z -component operators s_{1z} and s_{2z} are coupled to bath spins represented by σ_{nkz} . Here $n = 1, 2$ labels the baths and $k = 1, 2, 3, \dots, N_n$ labels the individual spins in the baths. For exchange symmetric case parameters of the two baths are taken to be identical.

In conclusion, exchange symmetry of a state, undergoing an external noise having the same symmetry, can be spontaneously broken as a result of decoherence due to local interactions. A natural question is the maximum probability of finding a symmetric possible final state as the system evolves in time. Even though we don't have a general analytical solution yet, our calculations show that total probability of conservation of exchange symmetry is $1/d$ at least for small systems. Evaluation of probabilities for possible broken symmetry states for higher d values will shed light on stability of superconducting state.

This work has been partially supported by the Scientific and Technological Research Council of Turkey (TUBITAK) under grant 107T530. Authors would like to thank the Institute of Theoretical and Applied Physics at Turunç where part of this research has been done.

References

- [1] P. W. Anderson, J. Phys. Chem. Solids 11 (1959) 26.
- [2] Z. Gedik, Solid State Commun. 124 (2002) 473.
- [3] B.R. Mottelson, The many-body problem, Ecole d'Eté de Physique Théorique Les Houches, Wiley, New York (1959).
- [4] I.O. Kulik, H. Boyaci, and Z. Gedik, Physica C 352 (2001) 46.
- [5] G. Karpat and Z. Gedik, Optics Commun. (2009) *in press*.
- [6] Ö. Bozat and Z. Gedik, Solid State Commun. 120 (2001) 487.
- [7] K. Kraus, States, Effects and Operations: Fundamental Notions of Quantum Theory, Springer-Verlag, Berlin, 1983.
- [8] D. Salgado, J. L. Sánchez-Gómez, Open Sys. and Information Dyn. 12 (2005) 5564.
- [9] M.D. Choi, Lin. Alg. and Appl. 10 (1975) 285.
- [10] Z. Gedik, Solid State Comm. 138 (2006) 82.