

# PLURIPOTENTIAL THEORY AND CONVEX BODIES: LARGE DEVIATION PRINCIPLE

TURGAY BAYRAKTAR, THOMAS BLOOM, NORMAN LEVENBERG,  
AND CHINH H. LU

ABSTRACT. We continue the study in [2] in the setting of weighted pluripotential theory arising from polynomials associated to a convex body  $P$  in  $(\mathbb{R}^+)^d$ . Our goal is to establish a large deviation principle in this setting specifying the rate function in terms of  $P$ -pluripotential-theoretic notions. As an important preliminary step, we first give an existence proof for the solution of a Monge-Ampère equation in an appropriate finite energy class. This is achieved using a variational approach.

## CONTENTS

1. Introduction	1
2. Monge-Ampère and $P$ -pluripotential theory	3
3. $P$ -pluripotential theory notions	13
4. Relation between $E^*$ and $J, J^Q$ functionals.	24
5. Large deviation.	30
References	37

## 1. INTRODUCTION

As in [2], we fix a convex body  $P \subset (\mathbb{R}^+)^d$  and we define the logarithmic indicator function

$$(1.1) \quad H_P(z) := \sup_{J \in P} \log |z^J| := \sup_{(j_1, \dots, j_d) \in P} \log[|z_1|^{j_1} \cdots |z_d|^{j_d}].$$

---

*Date:* February 10, 2019.

*2010 Mathematics Subject Classification.* 32U15, 32U20, 31C15.

*Key words and phrases.* convex body,  $P$ -extremal function, large deviation principle.

N. Levenberg is supported by Simons Foundation grant No. 354549.

We assume throughout that

$$(1.2) \quad \Sigma \subset kP \text{ for some } k \in \mathbb{Z}^+$$

where

$$\Sigma := \{(x_1, \dots, x_d) \in \mathbb{R}^d : 0 \leq x_i \leq 1, \sum_{j=1}^d x_j \leq 1\}.$$

Then

$$H_P(z) \geq \frac{1}{k} \max_{j=1, \dots, d} \log^+ |z_j|$$

where  $\log^+ |z_j| = \max[0, \log |z_j|]$ . We define

$$L_P = L_P(\mathbb{C}^d) := \{u \in PSH(\mathbb{C}^d) : u(z) - H_P(z) = O(1), |z| \rightarrow \infty\},$$

and

$$L_{P,+} = L_{P,+}(\mathbb{C}^d) = \{u \in L_P(\mathbb{C}^d) : u(z) \geq H_P(z) + C_u\}.$$

These are generalizations of the classical Lelong classes when  $P = \Sigma$ .

We define the finite-dimensional polynomial spaces

$$Poly(nP) := \{p(z) = \sum_{J \in nP \cap (\mathbb{Z}^+)^d} c_J z^J : c_J \in \mathbb{C}\}$$

for  $n = 1, 2, \dots$  where  $z^J = z_1^{j_1} \cdots z_d^{j_d}$  for  $J = (j_1, \dots, j_d)$ . For  $p \in Poly(nP)$ ,  $n \geq 1$  we have  $\frac{1}{n} \log |p| \in L_P$ ; also each  $u \in L_{P,+}(\mathbb{C}^d)$  is locally bounded in  $\mathbb{C}^d$ . For  $P = \Sigma$ , we write  $Poly(nP) = \mathcal{P}_n$ .

Given a compact set  $K \subset \mathbb{C}^d$ , one can define various pluripotential-theoretic notions associated to  $K$  related to  $L_P$  and the polynomial spaces  $Poly(nP)$ . Our goal in this paper is to prove some probabilistic properties of random point processes on  $K$  utilizing these notions and their weighted counterparts. We require an existence proof for the solution of a Monge-Ampère equation in an appropriate finite energy class; this is done in Theorem 2.8 using a variational approach and is of interest on its own. The third section recalls appropriate definitions and properties in  $P$ -pluripotential theory, mostly following [2]. As in [2], our spaces  $Poly(nP)$  do not necessarily arise as holomorphic sections of tensor powers of a line bundle. Subsection 3.3 includes a standard elementary probabilistic result on almost sure convergence of probability measures associated to random arrays on  $K$  to a  $P$ -pluripotential-theoretic equilibrium measure. Section 4 sets up the machinery for the more subtle large deviation principle (LDP), Theorem 5.1, for which we provide two proofs (analogous to those in [9]). As in [9], the first

proof was inspired by [6] and the second proof was utilized by Berman in [5]. The reader will find far-reaching applications and interpretations of LDP's in the appropriate settings of holomorphic line bundles over a compact, complex manifold in [5]. In particular, the case where  $P$  is a convex integral polytope (vertices in  $\mathbb{Z}^d$ ) which is the moment polytope for a toric manifold ( $P$  is Delzant) is covered in [5].

## 2. MONGE-AMPÈRE AND $P$ -PLURIPOTENTIAL THEORY

**2.1. Monge-Ampère equations with prescribed singularity.** In this section,  $(X, \omega)$  is a compact Kähler manifold of dimension  $d$ .

**2.1.1. Quasi-plurisubharmonic functions.** A function  $u : X \rightarrow \mathbb{R} \cup \{-\infty\}$  is called quasi-plurisubharmonic (quasi-psh) if locally  $u = \rho + \varphi$ , where  $\varphi$  is plurisubharmonic and  $\rho$  is smooth.

We let  $PSH(X, \omega)$  denote the set of  $\omega$ -psh functions, i.e. quasi-psh functions  $u$  such that  $\omega_u := \omega + dd^c u \geq 0$  in the sense of currents on  $X$ .

Given  $u, v \in PSH(X, \omega)$  we say that  $u$  is more singular than  $v$  (and we write  $u \prec v$ ) if  $u \leq v + C$  on  $X$ , for some constant  $C$ . We say that  $u$  has the same singularity as  $v$  (and we write  $u \simeq v$ ) if  $u \prec v$  and  $v \prec u$ .

Given  $\phi \in PSH(X, \omega)$ , we let  $PSH(X, \omega, \phi)$  denote the set of  $\omega$ -psh functions  $u$  which are more singular than  $\phi$ .

**2.1.2. Nonpluripolar Monge-Ampère measure.** For bounded  $\omega$ -psh functions  $u_1, \dots, u_d$ , the Monge-Ampère product  $(\omega + dd^c u_1) \wedge \dots \wedge (\omega + dd^c u_d)$  is well-defined as a positive Radon measure on  $X$  (see [14], [3]). For general  $\omega$ -psh functions  $u_1, \dots, u_d$ , the sequence of positive measures

$$\mathbf{1}_{\cap\{u_j > -k\}}(\omega + dd^c \max(u_1, -k)) \wedge \dots \wedge (\omega + dd^c \max(u_d, -k))$$

is non-decreasing in  $k$  and the limiting measure, which is called the nonpluripolar product of  $\omega_{u_1}, \dots, \omega_{u_d}$ , is denoted by

$$\omega_{u_1} \wedge \dots \wedge \omega_{u_d}.$$

When  $u_1 = \dots = u_d = u$  we write  $\omega_u^d := \omega_u \wedge \dots \wedge \omega_u$ . Note that by definition  $\int_X \omega_{u_1} \wedge \dots \wedge \omega_{u_d} \leq \int_X \omega^d$ .

It was proved in [20, Theorem 1.2] and [11, Theorem 1.1] that the total mass of nonpluripolar Monge-Ampère products is decreasing with respect to singularity type. More precisely,

**Theorem 2.1.** *Let  $\omega_1, \dots, \omega_d$  be Kähler forms on  $X$ . If  $u_j \leq v_j$ ,  $j = 1, \dots, d$ , are  $\omega_j$ -psh functions then*

$$\int_X (\omega_1 + dd^c u_1) \wedge \dots \wedge (\omega_d + dd^c u_d) \leq \int_X (\omega_1 + dd^c v_1) \wedge \dots \wedge (\omega_d + dd^c v_d).$$

As noted above, for a general  $\omega$ -psh function  $u$  we have the estimate  $\int_X \omega_u^d \leq \int_X \omega^d$ . Following [15] we let  $\mathcal{E}(X, \omega)$  denote the set of all  $\omega$ -psh functions with maximal total mass, i.e.

$$\mathcal{E}(X, \omega) := \left\{ u \in PSH(X, \omega) : \int_X \omega_u^d = \int_X \omega^d \right\}.$$

Given  $\phi \in PSH(X, \omega)$ , we define

$$\mathcal{E}(X, \omega, \phi) := \left\{ u \in PSH(X, \omega, \phi) : \int_X \omega_u^d = \int_X \omega_\phi^d \right\}.$$

**Proposition 2.2.** *Let  $\phi \in PSH(X, \omega)$ . The following are equivalent :*

- (1)  $\mathcal{E}(X, \omega, \phi) \cap \mathcal{E}(X, \omega) \neq \emptyset$ ;
- (2)  $\phi \in \mathcal{E}(X, \omega)$ ;
- (3)  $\mathcal{E}(X, \omega, \phi) \subset \mathcal{E}(X, \omega)$ .

*Proof.* We first prove (1)  $\implies$  (2). If  $u \in \mathcal{E}(X, \omega, \phi) \cap \mathcal{E}(X, \omega)$  then  $\int_X \omega_u^d = \int_X \omega^d$ . On the other hand, since  $u$  is more singular than  $\phi$ , Theorem 2.1 ensures that

$$\int_X \omega^d = \int_X \omega_u^d \leq \int_X \omega_\phi^d \leq \int_X \omega^d,$$

hence equality holds, proving that  $\phi \in \mathcal{E}(X, \omega)$ .

Now we prove (2)  $\implies$  (3). If  $\phi \in \mathcal{E}(X, \omega)$  and  $u \in \mathcal{E}(X, \omega, \phi)$  then

$$\int_X \omega_u^d = \int_X \omega_\phi^d = \int_X \omega^d,$$

hence  $u \in \mathcal{E}(X, \omega)$ .

Finally (3)  $\implies$  (1) is obvious.  $\square$

**Proposition 2.3.** *Assume that  $\phi_j \in PSH(X, \omega_j)$ ,  $j = 1, \dots, d$  with  $\int_X (\omega_j + dd^c \phi_j)^d > 0$ . If  $u_j \in \mathcal{E}(X, \omega_j, \phi_j)$ ,  $j = 1, \dots, d$ , then*

$$\int_X (\omega_1 + dd^c u_1) \wedge \dots \wedge (\omega_d + dd^c u_d) = \int_X (\omega_1 + dd^c \phi_1) \wedge \dots \wedge (\omega_d + dd^c \phi_d).$$

*Proof.* Theorem 2.1 gives one inequality. The other one follows from [11, Proposition 3.1 and Theorem 3.14].  $\square$

2.1.3. *Model potentials.* For a function  $f : X \rightarrow \mathbb{R} \cup \{-\infty\}$ , we let  $f^*$  denote its uppersemicontinuous (usc) regularization, i.e.

$$f^*(x) := \limsup_{X \ni y \rightarrow x} f(y).$$

Given  $\phi \in PSH(X, \omega)$ , following J. Ross and D. Witt Nyström [18], we define

$$P_\omega[\phi] := \left( \lim_{t \rightarrow +\infty} P_\omega(\min(\phi + t, 0)) \right)^*.$$

Here, for a function  $f$ ,  $P_\omega(f)$  is defined as

$$P_\omega(f) := (x \mapsto \sup\{u(x) : u \in PSH(X, \omega), u \leq f\})^*.$$

It was shown in [11, Theorem 3.8] that the nonpluripolar Monge-Ampère measure of  $P_\omega[\phi]$  is dominated by Lebesgue measure:

$$(2.1) \quad (\omega + dd^c P_\omega[\phi])^d \leq \mathbf{1}_{\{P_\omega[\phi]=0\}} \omega^d \leq \omega^d.$$

This fact plays a crucial role in solving the complex Monge-Ampère equation. For the reader's convenience, we note that in the notation of [11] (on the left)

$$P_{[\omega, \phi]}(0) = P_\omega[\phi].$$

**Definition 2.4.** A function  $\phi \in PSH(X, \omega)$  is called a model potential if  $\int_X \omega_\phi^d > 0$  and  $P_\omega[\phi] = \phi$ . A function  $u \in PSH(X, \omega)$  has model type singularity if  $u$  has the same singularity as  $P_\omega[u]$ ; i.e.,  $u - P_\omega[u]$  is bounded on  $X$ .

There are plenty of model potentials. If  $\varphi \in PSH(X, \omega)$  with  $\int_X \omega_\varphi^d > 0$  then, by [11, Theorem 3.12],  $P_\omega[\varphi]$  is a model potential. In particular, if  $\int_X \omega_\varphi^d = \int_X \omega^d$  (i.e.  $\varphi \in \mathcal{E}(X, \omega)$ ) then  $P_\omega[\varphi] = 0$ .

We will use the following property of model potentials proved in [11, Theorem 3.12]: if  $\phi$  is a model potential then

$$(2.2) \quad u \in PSH(X, \omega, \phi) \implies u - \sup_X u \leq \phi.$$

In the sequel we always assume that  $\phi$  has *model type singularity* and *small unbounded locus*; i.e.,  $\phi$  is locally bounded outside a closed complete pluripolar set, allowing us to use the variational approach of [7] as explained in [11].

2.1.4. *The variational approach.* We call a measure which puts no mass on pluripolar sets a *nonpluripolar measure*. For a positive nonpluripolar measure  $\mu$  on  $X$  we let  $L_\mu$  denote the following linear functional on  $PSH(X, \omega, \phi)$ :

$$L_\mu(u) := \int_X (u - \phi) d\mu.$$

For  $u \in PSH(X, \omega)$  with  $u \simeq \phi$ , we define the Monge-Ampère energy

$$(2.3) \quad \mathbf{E}_\phi(u) := \frac{1}{(d+1)} \sum_{k=0}^d \int_X (u - \phi) \omega_u^k \wedge \omega_\phi^{d-k}.$$

It was shown in [11, Theorem 4.10] (by adapting the arguments of [7]) that  $\mathbf{E}_\phi$  is non-decreasing and concave along affine curves, giving rise to its trivial extension to  $PSH(X, \omega, \phi)$ .

We define

$$(2.4) \quad \mathcal{E}^1(X, \omega, \phi) := \{u \in PSH(X, \omega, \phi) : \mathbf{E}_\phi(u) > -\infty\}.$$

The following criterion was proved in [11, Theorem 4.13]:

**Proposition 2.5.** *Let  $u \in PSH(X, \omega, \phi)$ . Then  $u \in \mathcal{E}^1(X, \omega, \phi)$  iff  $u \in \mathcal{E}(X, \omega, \phi)$  and  $\int_X (u - \phi) \omega_u^d > -\infty$ .*

**Lemma 2.6.** *If  $E$  is pluripolar then there exists  $u \in \mathcal{E}^1(X, \omega, \phi)$  such that  $E \subset \{u = -\infty\}$ .*

*Proof.* Without loss of generality we can assume that  $\phi$  is a model potential. Then (2.1) gives  $\int_X |\phi| \omega_\phi^d = 0$ . It follows from [7, Corollary 2.11] that there exists  $v \in \mathcal{E}^1(X, \omega, 0)$ ,  $v \leq 0$ , such that  $E \subset \{v = -\infty\}$ . Set  $u := P_\omega(\min(v, \phi))$ . Then  $E \subset \{u = -\infty\}$  and we claim that  $u \in \mathcal{E}^1(X, \omega, \phi)$ . For each  $j \in \mathbb{N}$  we set  $v_j := \max(v, -j)$  and  $u_j := P_\omega(\min(v_j, \phi))$ . Then  $u_j$  decreases to  $u$  and  $u_j \simeq \phi$ . Using [11, Theorem 4.10 and Lemma 4.15] it suffices to check that  $\{\int_X |u_j - \phi| \omega_{u_j}^d\}$  is uniformly bounded. It follows from [11, Lemma 3.7] that

$$\begin{aligned} \int_X |u_j - \phi| \omega_{u_j}^d &\leq \int_X |u_j| \omega_{u_j}^d \leq \int_X |v_j| \omega_{v_j}^d + \int_X |\phi| \omega_\phi^d \\ &= \int_X |v_j| \omega_{v_j}^d. \end{aligned}$$

The fact that  $\int_X |v_j| \omega_{v_j}^d$  is uniformly bounded follows from [15, Corollary 2.4] since  $v \in \mathcal{E}^1(X, \omega, 0)$ . This concludes the proof.  $\square$

**Lemma 2.7.** *Assume that  $\mathcal{E}^1(X, \omega, \phi) \subset L^1(X, \mu)$ . Then, for each  $C > 0$ ,  $L_\mu$  is bounded on*

$$E_C := \{u \in PSH(X, \omega, \phi) : \sup_X u \leq 0 \text{ and } \mathbf{E}_\phi(u) \geq -C\}.$$

*Proof.* By concavity of  $\mathbf{E}_\phi$  the set  $E_C$  is convex. We now show that  $E_C$  is compact in the  $L^1(X, \omega^d)$  topology. Let  $\{u_j\}$  be a sequence in  $E_C$ . We claim that  $\{\sup_X u_j\}$  is bounded. Indeed, by [11, Theorem 4.10]

$$\begin{aligned} \mathbf{E}_\phi(u_j) &\leq \int_X (u_j - \phi) \omega_\phi^d \\ &\leq (\sup_X u_j) \int_X \omega_\phi^d + \int_X (u_j - \sup_X u_j - \phi) \omega_\phi^d. \end{aligned}$$

It follows from (2.2) that  $u_j - \sup_X u_j \leq P_\omega[\phi] \leq \phi + C_0$ , where  $C_0$  is a constant. The boundedness of  $\{\sup_X u_j\}$  then follows from that of  $\{\mathbf{E}_\phi(u_j)\}$  and the above estimate. This proves the claim.

A subsequence of  $\{u_j\}$ , still denoted by  $\{u_j\}$ , converges in  $L^1(X, \omega^d)$  to  $u \in PSH(X, \omega)$  with  $\sup_X u \leq 0$ . Since  $u_j - \sup_X u_j \leq \phi + C_0$ , we have  $u - \sup_X u \leq \phi + C_0$ . This proves that  $u \in PSH(X, \omega, \phi)$ . The upper semicontinuity of  $\mathbf{E}_\phi$  (see [11, Proposition 4.19]) ensures that  $\mathbf{E}_\phi(u) \geq -C$ , hence  $u \in E_C$ . This proves that  $E_C$  is compact in the  $L^1(X, \omega^d)$  topology.

The result then follows from [7, Proposition 3.4].  $\square$

The goal of this section is to prove the following result:

**Theorem 2.8.** *Assume that  $\mu$  is a nonpluripolar positive measure on  $X$  such that  $\mu(X) = \int_X \omega_\phi^d$ . The following are equivalent*

- (1)  $\mu$  has finite energy, i.e.,  $L_\mu$  is finite on  $\mathcal{E}^1(X, \omega, \phi)$ ;
- (2) there exists  $u \in \mathcal{E}^1(X, \omega, \phi)$  such that  $\omega_u^d = \mu$ ;
- (3) there exists a unique  $u \in \mathcal{E}^1(X, \omega, \phi)$  such that

$$F_\mu(u) = \max_{v \in \mathcal{E}^1(X, \omega, \phi)} F_\mu(v) < +\infty$$

where  $F_\mu = \mathbf{E}_\phi - L_\mu$ .

**Remark 2.9.** It was shown in [11, Theorem 4.28] that a unique (normalized) solution  $u$  in  $\mathcal{E}(X, \omega, \phi)$  always exists (without the finite energy assumption on  $\mu$ ). But that proof does not give a solution in  $\mathcal{E}^1(X, \omega, \phi)$ . Below, we will follow the proof of [11, Theorem 4.28] and use the finite energy condition,  $\mathcal{E}^1(X, \omega, \phi) \subset L^1(X, \mu)$ , to prove that  $u$  belongs to  $\mathcal{E}^1(X, \omega, \phi)$ .

**Lemma 2.10.** *Assume that  $\mathcal{E}^1(X, \omega, \phi) \subset L^1(X, \mu)$ . Then there exists a positive constant  $C$  such that, for all  $u \in \mathcal{E}^1(X, \omega, \phi)$  with  $\sup_X u = 0$ ,*

$$(2.5) \quad L_\mu(u) \geq -C(1 + |\mathbf{E}_\phi(u)|^{1/2}).$$

The proof below uses ideas in [15, 7].

*Proof.* Since  $\phi$  has model type singularity, it follows from [11, Theorem 4.10] that  $\mathbf{E}_\phi - \mathbf{E}_{P_\omega[\phi]}$  is bounded. Without loss of generality we can assume in this proof that  $\phi = P_\omega[\phi]$ . Fix  $u \in \mathcal{E}^1(X, \omega, \phi)$  such that  $\sup_X u = 0$  and  $|\mathbf{E}_\phi(u)| > 1$ . Then, by [11, Theorem 3.12],  $u \leq \phi$ . Set  $a = |\mathbf{E}_\phi(u)|^{-1/2} \in (0, 1)$ , and  $v := au + (1 - a)\phi \in \mathcal{E}^1(X, \omega, \phi)$ . We estimate  $\mathbf{E}_\phi(v)$  as follows

$$\begin{aligned} (d+1)\mathbf{E}_\phi(v) &= a \sum_{k=0}^d \int_X (u - \phi) \omega_u^k \wedge \omega_\phi^{d-k} \\ &= a \sum_{k=0}^d \int_X (u - \phi) (a\omega_u + (1-a)\omega_\phi)^k \wedge \omega_\phi^{d-k} \\ &\geq C(d)a \int_X (u - \phi) \omega_\phi^d + C(d)a^2 \sum_{k=0}^d \int_X (u - \phi) \omega_u^k \wedge \omega_\phi^d, \end{aligned}$$

where  $C(d)$  is a positive constant which only depends on  $d$ . It follows from  $\phi = P_\omega[\phi]$  and [11, Theorem 3.8] that  $\omega_\phi^d \leq \omega^d$  (recall (2.1)). This together with [14, Proposition 2.7] give

$$\int_X (u - \phi) \omega_\phi^d \geq -C_1,$$

for a uniform constant  $C_1$ . Therefore,

$$(d+1)\mathbf{E}_\phi(v) \geq -C_1 C(d)a + C_2 a^2 \mathbf{E}_\phi(u) \geq -C_3.$$

It thus follows from Lemma 2.7 that  $L_\mu(v) \geq -C_4$  for a uniform constant  $C_4 > 0$ . Thus

$$\int_X (u - \phi) d\mu \geq -C_4/a,$$

which gives (2.5). □

We are now ready to prove Theorem 2.8.



*Proof of Theorem 2.8.* Without loss of generality we can assume that  $\phi$  is a model potential. We first prove (1)  $\implies$  (2). We write  $\mu = f\nu$ , where  $\nu$  is a nonpluripolar positive measure satisfying, for all Borel subsets  $B \subset X$ ,

$$\nu(B) \leq A \text{Cap}_\phi(B),$$

for some positive constant  $A$ , and  $0 \leq f \in L^1(X, \nu)$  (cf., [11, Lemma 4.26]). Here  $\text{Cap}_\phi$  is defined as

$$\text{Cap}_\phi(B) := \sup \left\{ \int_B \omega_u^d : u \in PSH(X, \omega), \phi - 1 \leq u \leq \phi \right\}.$$

Set, for  $k \in \mathbb{N}$ ,  $\mu_k := c_k \min(f, k)\nu$  where  $c_k > 0$  is chosen so that  $\mu_k(X) = \int_X \omega_\phi^d$ ; this is needed in order to solve the Monge-Ampère equation in the class  $\mathcal{E}^1(X, \omega, \phi)$ . For  $k$  large enough,  $1 \leq c_k \leq 2$  and  $c_k \rightarrow 1$  as  $k \rightarrow +\infty$ . It follows from [11, Theorem 4.25] that there exists  $u_j \in \mathcal{E}^1(X, \omega, \phi)$ ,  $\sup_X u_j = 0$ , such that  $\omega_{u_j}^d = \mu_j$ ; by [11, Theorem 3.12],  $u_j \leq \phi$ . A subsequence of  $\{u_j\}$  which, by abuse of notation, will be denoted by  $\{u_j\}$ , converges in  $L^1(X, \mu)$  to  $u \in PSH(X, \omega)$  with  $u \leq \phi$ . Define  $v_k := (\sup_{j \geq k} u_j)^*$ . Then  $v_k \searrow u$  and  $\sup_X v_k = 0$ . It follows from (2.5) and [11, Theorem 4.10] that

$$\begin{aligned} |\mathbf{E}_\phi(u_j)| &\leq \int_X |u_j - \phi| \omega_{u_j}^d \leq 2 \int_X |u_j - \phi| d\mu \\ &\leq 2C(1 + |\mathbf{E}_\phi(u_j)|^{1/2}). \end{aligned}$$

Therefore  $\{|\mathbf{E}_\phi(u_j)|\}$  is bounded, hence so is  $\{|\mathbf{E}_\phi(v_j)|\}$  since  $\mathbf{E}_\phi$  is non-decreasing. It then follows from [11, Lemma 4.15] that  $u \in \mathcal{E}^1(X, \omega, \phi)$ .

Now, repeating the arguments of [11, Theorem 4.28] we can show that  $\omega_u^d = \mu$ , finishing the proof of (1)  $\implies$  (2).

We next prove (2)  $\implies$  (3). Assume that  $\mu = \omega_u^d$  for some  $u \in \mathcal{E}^1(X, \omega, \phi)$ . For all  $v \in \mathcal{E}^1(X, \omega, \phi)$ , by [11, Theorem 4.10] and Proposition 2.5 we have

$$\begin{aligned} L_\mu(v) &= \int_X (v - \phi) \omega_u^d \\ &= \int_X (v - u) \omega_u^d + \int_X (u - \phi) \omega_u^d \\ &\geq \mathbf{E}_\phi(v) - \mathbf{E}_\phi(u) + \int_X (u - \phi) \omega_u^d > -\infty. \end{aligned}$$

Hence  $L_\mu$  is finite on  $\mathcal{E}^1(X, \omega, \phi)$ . Now, for all  $v \in \mathcal{E}^1(X, \omega, \phi)$ , by [11, Theorem 4.10] we have

$$F_\mu(v) - F_\mu(u) = \mathbf{E}_\phi(v) - \mathbf{E}_\phi(u) - \int_X (v - u)\omega_u^d \leq 0.$$

This gives (3). Finally, (3)  $\implies$  (1) is obvious.  $\square$

## 2.2. Monge-Ampère equations on $\mathbb{C}^d$ with prescribed growth.

As in the introduction we let  $P$  be a convex body contained in  $(\mathbb{R}^+)^d$  and fix  $r > 0$  such that  $P \subset r\Sigma$ . We assume (1.2); i.e.,  $\Sigma \subset kP$  for some  $k \in \mathbb{Z}^+$ . This ensures that  $H_P$  in (1.1) is locally bounded on  $\mathbb{C}^d$  (and of course  $H_P \in L_P^+(\mathbb{C}^d)$ ). Let  $u \in L_P(\mathbb{C}^d)$  and define

$$(2.6) \quad \tilde{u}(z) := u(z) - \frac{r}{2} \log(1 + |z|^2), z \in \mathbb{C}^d.$$

Consider the projective space  $\mathbb{P}^d$  equipped with the Kähler metric  $\omega := r\omega_{FS}$ , where

$$\omega_{FS} = dd^c \frac{1}{2} \log(1 + |z|^2)$$

on  $\mathbb{C}^d$ . Then  $\tilde{u}$  is bounded from above on  $\mathbb{C}^d$ . It thus can be extended to  $\mathbb{P}^d$  as a function in  $PSH(\mathbb{P}^d, \omega)$ .

For a plurisubharmonic function  $u$  on  $\mathbb{C}^d$ , we let  $(dd^c u)^d$  denotes its nonpluripolar Monge-Ampère measure; i.e.,  $(dd^c u)^d$  is the increasing limit of the sequence of measures  $\mathbf{1}_{\{u > -k\}}(dd^c \max(u, -k))^d$ . Then

$$\omega_{\tilde{u}}^d = (\omega + dd^c \tilde{u})^d = (dd^c u)^d \text{ on } \mathbb{C}^d.$$

If  $u \in L_P(\mathbb{C}^d)$  then

$$\int_{\mathbb{C}^d} (dd^c u)^d \leq \int_{\mathbb{C}^d} (dd^c H_P)^d = d! \text{Vol}(P) =: \gamma_d = \gamma_d(P)$$

(cf., equation (2.4) in [2]). We define

$$\mathcal{E}_P(\mathbb{C}^d) := \left\{ u \in L_P(\mathbb{C}^d) : \int_{\mathbb{C}^d} (dd^c u)^d = \gamma_d \right\}.$$

By the construction in (2.6) we have that  $\tilde{H}_P \in PSH(\mathbb{P}^d, \omega)$ . We define

$$\tilde{\Phi}_P := P_\omega[\tilde{H}_P].$$

The key point here, which follows from [12, Theorem 7.2], is that  $\tilde{H}_P$  has model type singularity (recall Definition 2.4) and hence the same

singularity as  $\tilde{\Phi}_P$ . Defining  $\Phi_P$  on  $\mathbb{C}^d$  using (2.6); i.e., for  $z \in \mathbb{C}^d$ ,

$$\Phi_P(z) = \tilde{\Phi}_P(z) + \frac{r}{2} \log(1 + |z|^2),$$

we thus have  $\Phi_P \in L_{P,+}(\mathbb{C}^d)$ . The advantage of using  $\Phi_P$  is that, by (2.1),  $(dd^c\Phi_P)^d \leq \omega^d$  on  $\mathbb{C}^d$ . Note that  $L_{P,+}(\mathbb{C}^d) \subset \mathcal{E}_P(\mathbb{C}^d)$ . For  $u, v \in L_P^+(\mathbb{C}^d)$  we define

$$(2.7) \quad E_v(u) := \frac{1}{(d+1)} \sum_{j=0}^d \int_{\mathbb{C}^d} (u-v)(dd^c u)^j \wedge (dd^c v)^{d-j}.$$

The corresponding global energy (see (2.3)) is defined as

$$\mathbf{E}_{\tilde{v}}(\tilde{u}) := \frac{1}{(d+1)} \sum_{j=0}^d \int_{\mathbb{P}^d} (\tilde{u} - \tilde{v})(\omega + dd^c \tilde{u})^j \wedge (\omega + dd^c \tilde{v})^{d-j}.$$

Then  $E_v$  is non-decreasing and concave along affine curves in  $L_{P,+}(\mathbb{C}^d)$ . We extend  $E_v$  to  $L_P(\mathbb{C}^d)$  in an obvious way. Note that  $E_v$  may take the value  $-\infty$ . We define

$$\mathcal{E}_P^1(\mathbb{C}^d) := \{u \in L_P(\mathbb{C}^d) : E_{H_P}(u) > -\infty\}.$$

We observe that in the above definition we can replace  $E_{H_P}$  by  $E_{\Phi_P}$ , since for  $u \in L_{P,+}(\mathbb{C}^d)$ , by the cocycle property (cf. Proposition 3.3 [2]),

$$E_{H_P}(u) - E_{H_P}(\Phi_P) = E_{\Phi_P}(u).$$

We thus have the following important identification (see (2.4)):

$$(2.8) \quad u \in \mathcal{E}_P^1(\mathbb{C}^d) \iff \tilde{u} \in \mathcal{E}^1(\mathbb{P}^d, \omega, \tilde{\Phi}_P).$$

We then have the following local version of Proposition 2.5:

**Proposition 2.11.** *Let  $u \in L_P(\mathbb{C}^d)$ . Then  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$  iff  $u \in \mathcal{E}_P(\mathbb{C}^d)$  and  $\int_{\mathbb{C}^d} (u - H_P)(dd^c u)^d > -\infty$ . In particular, if  $\text{supp}(dd^c u)^d$  is compact,  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$  iff  $\int_{\mathbb{C}^d} (dd^c u)^d = \gamma_d$  and  $\int_{\mathbb{C}^d} u(dd^c u)^d > -\infty$ .*

*Proof.* Since  $\tilde{H}_P \simeq \tilde{\Phi}_P$ ,

$$\int_{\mathbb{P}^d} (\tilde{u} - \tilde{H}_P)\omega_{\tilde{u}}^d > -\infty \text{ iff } \int_{\mathbb{P}^d} (\tilde{u} - \tilde{\Phi}_P)\omega_{\tilde{u}}^d > -\infty$$

where  $\tilde{u} \in PSH(\mathbb{P}^d, \omega)$  and  $u$  are related by (2.6). Moreover,  $\Phi_P \in L_{P,+}(\mathbb{C}^d)$  implies  $u \leq \Phi_P + c$  so that  $\tilde{u} \in PSH(\mathbb{P}^d, \omega, \tilde{\Phi}_P)$ . But

$$\int_{\mathbb{P}^d} (\tilde{u} - \tilde{H}_P)\omega_{\tilde{u}}^d = \int_{\mathbb{C}^d} (u - H_P)(dd^c u)^d$$

and the result follows from (2.8) by applying Proposition 2.5 to  $\tilde{u}$ . For the last statement, note that for general  $u \in L_P(\mathbb{C}^d)$  we may have  $\int_{\mathbb{C}^d} H_P(dd^c u)^d = +\infty$ , but if  $(dd^c u)^d$  has compact support then  $\int_{\mathbb{C}^d} H_P(dd^c u)^d$  is finite.  $\square$

Note that Theorem 2.1 and Proposition 2.3 give the following result:

**Theorem 2.12.** *Let  $u_1, \dots, u_d$  be functions in  $\mathcal{E}_P(\mathbb{C}^d)$ . Then*

$$\int_{\mathbb{C}^d} dd^c u_1 \wedge \dots \wedge dd^c u_d = \gamma_d.$$

For  $u_1, \dots, u_n \in L_{P,+}(\mathbb{C}^d)$  Theorem 2.12 was proved in [1, Proposition 2.7].

Having the correspondence (2.8) we can state a local version of Theorem 2.8; this will be used in the sequel. Let  $\mathcal{M}_P(\mathbb{C}^d)$  denote the set of all positive Borel measures  $\mu$  on  $\mathbb{C}^d$  with  $\mu(\mathbb{C}^d) = d! \text{Vol}(P) = \gamma_d$ .

**Theorem 2.13.** *Assume that  $\mu \in \mathcal{M}_P(\mathbb{C}^d)$  is a positive nonpluripolar Borel measure. The following are equivalent*

- (1)  $\mathcal{E}_P^1(\mathbb{C}^d) \subset L^1(\mathbb{C}^d, \mu)$ ;
- (2) there exists  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$  such that  $(dd^c u)^d = \mu$ ;
- (3) there exists  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$  such that

$$\mathcal{F}_\mu(u) = \max_{v \in \mathcal{E}_P^1(\mathbb{C}^d)} \mathcal{F}_\mu(v) < +\infty.$$

A priori the functional  $\mathcal{F}_\mu$  is defined for  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$  by

$$\mathcal{F}_{\mu, \Phi_P}(u) := E_{\Phi_P}(u) - \int_{\mathbb{C}^d} (u - \Phi_P) d\mu.$$

However, using this notation, since

$$\mathcal{F}_{\mu, \Phi_P}(u) - \mathcal{F}_{\mu, H_P}(u) = \mathcal{F}_{\mu, \Phi_P}(H_P),$$

in statement (3) of Theorem 2.13 we can take either of the two definitions  $\mathcal{F}_{\mu, \Phi_P}$  or  $\mathcal{F}_{\mu, H_P}$  for  $\mathcal{F}_\mu$ .

**Remark 2.14.** If  $\mu$  has compact support in  $\mathbb{C}^d$  then  $\int_{\mathbb{C}^d} \Phi_P d\mu$  and  $\int_{\mathbb{C}^d} H_P d\mu$  are finite. Therefore, the functional  $\mathcal{F}_\mu$  can be replaced by

$$u \mapsto E_{H_P}(u) - \int_{\mathbb{C}^d} u d\mu.$$

Using the remark, for  $\mu \in \mathcal{M}_P(\mathbb{C}^d)$  with compact support, it is natural to define the Legendre-type transform of  $E_{H_P}$ :

$$(2.9) \quad E^*(\mu) := \sup_{u \in \mathcal{E}_P^1(\mathbb{C}^d)} [E_{H_P}(u) - \int_{\mathbb{C}^d} u d\mu].$$

This functional, which will appear in the rate function for our LDP, will be given a more concrete interpretation using  $P$ -pluripotential theory in section 4; cf., equation (4.18).

Finally, for future use, we record the following consequence of Lemma 2.6 and the correspondence (2.8).

**Lemma 2.15.** *If  $E \subset \mathbb{C}^d$  is pluripolar then there exists  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$  such that  $E \subset \{u = -\infty\}$ .*

### 3. $P$ -PLURIPOTENTIAL THEORY NOTIONS

Given  $E \subset \mathbb{C}^d$ , the  $P$ -extremal function of  $E$  is

$$V_{P,E}^*(z) := \limsup_{\zeta \rightarrow z} V_{P,E}(\zeta)$$

where

$$V_{P,E}(z) := \sup\{u(z) : u \in L_P(\mathbb{C}^d), u \leq 0 \text{ on } E\}.$$

For  $K \subset \mathbb{C}^d$  compact,  $w : K \rightarrow \mathbb{R}^+$  is an admissible weight function on  $K$  if  $w \geq 0$  is an uppersemicontinuous function with  $\{z \in K : w(z) > 0\}$  nonpluripolar. Setting  $Q := -\log w$ , we write  $Q \in \mathcal{A}(K)$  and define the *weighted  $P$ -extremal function*

$$V_{P,K,Q}^*(z) := \limsup_{\zeta \rightarrow z} V_{P,K,Q}(\zeta)$$

where

$$V_{P,K,Q}(z) := \sup\{u(z) : u \in L_P(\mathbb{C}^d), u \leq Q \text{ on } K\}.$$

If  $Q = 0$  we write  $V_{P,K,Q} = V_{P,K}$ , consistent with the previous notation. For  $P = \Sigma$ ,

$$V_{\Sigma,K,Q}(z) = V_{K,Q}(z) := \sup\{u(z) : u \in L(\mathbb{C}^d), u \leq Q \text{ on } K\}$$

is the usual weighed extremal function as in Appendix B of [19].

We write (omitting the dependence on  $P$ )

$$\mu_{K,Q} := (dd^c V_{P,K,Q}^*)^d \text{ and } \mu_K := (dd^c V_{P,K}^*)^d$$

for the Monge-Ampère measures of  $V_{P,K,Q}^*$  and  $V_{P,K}^*$  (the latter if  $K$  is not pluripolar). Proposition 2.5 of [2] states that

$$\text{supp}(\mu_{K,Q}) \subset \{z \in K : V_{P,K,Q}^*(z) \geq Q(z)\}$$

and  $V_{P,K,Q}^* = Q$  q.e. on  $\text{supp}(\mu_{K,Q})$ , i.e., off of a pluripolar set.

**3.1. Energy.** We recall some results and definitions from [2]. For  $u, v \in L_{P,+}(\mathbb{C}^d)$ , we define the *mutual energy*

$$\mathcal{E}(u, v) := \int_{\mathbb{C}^d} (u - v) \sum_{j=0}^d (dd^c u)^j \wedge (dd^c v)^{d-j}.$$

For simplicity, when  $v = H_P$ , we denote the associated (normalized) energy functional by  $E$ :

$$E(u) := E_{H_P}(u) = \frac{1}{d+1} \sum_{j=0}^d \int_{\mathbb{C}^d} (u - H_P) dd^c u^j \wedge (dd^c H_P)^{d-j}$$

(recall (2.7)).

For  $u, u', v \in L_{P,+}(\mathbb{C}^d)$ , and for  $0 \leq t \leq 1$ , we define

$$f(t) := \mathcal{E}(u + t(u' - u), v),$$

From Proposition 3.1 in [2],  $f'(t)$  exists for  $0 \leq t \leq 1$  and

$$f'(t) = (d+1) \int_{\mathbb{C}^d} (u' - u) (dd^c(u + t(u' - u)))^d$$

Hence, taking  $v = H_P$ , we have, for  $F(t) := E(u + t(u' - u))$ , that

$$F'(t) = \int_{\mathbb{C}^d} (u' - u) (dd^c(u + t(u' - u)))^d.$$

Thus  $F'(0) = \int_{\mathbb{C}^d} (u' - u) (dd^c u)^d$  and we write

$$(3.1) \quad \langle E'(u), u' - u \rangle := \int_{\mathbb{C}^d} (u' - u) (dd^c u)^d.$$

We need some applications of a global domination principle. The following version, sufficient for our purposes, follows from [11], Corollary 3.10 (see also Corollary A.2 of [8]).

**Proposition 3.1.** *Let  $u \in L_P(\mathbb{C}^d)$  and  $v \in \mathcal{E}_P(\mathbb{C}^d)$  with  $u \leq v$  a.e.  $(dd^c v)^d$ . Then  $u \leq v$  in  $\mathbb{C}^d$ .*

This will be used to prove an approximation result, Proposition 3.3, which itself will be essential in the sequel. First we need a lemma.

**Lemma 3.2.** *Assume that  $\varphi \leq u, v \leq H_P$  are functions in  $\mathcal{E}_P^1(\mathbb{C}^d)$ . Then for all  $t > 0$ ,*

$$\int_{\{u \leq H_P - 2t\}} (H_P - u)(dd^c v)^d \leq 2^{d+1} \int_{\{\varphi \leq H_P - t\}} (H_P - \varphi)(dd^c \varphi)^d.$$

*In particular, the left hand side converges to 0 as  $t \rightarrow +\infty$  uniformly in  $u, v$ .*

*Proof.* For  $s > 0$ , we have the following inclusions of sets:

$$(u \leq H_P - 2s) \subset \left( \varphi \leq \frac{v + H_P}{2} - s \right) \subset (\varphi \leq H_P - s).$$

We first note that the left hand side in the lemma is equal to

$$(3.2) \quad \int_{\{u \leq H_P - 2t\}} (H_P - u)(dd^c v)^d \\ = 2t \int_{\{u \leq H_P - 2t\}} (dd^c v)^d + \int_{2t}^{\infty} \left( \int_{\{u \leq H_P - s\}} (dd^c v)^d \right) ds.$$

We claim that, for all  $s > 0$ ,

$$(3.3) \quad \int_{\{u \leq H_P - 2s\}} (dd^c v)^d \leq 2^d \int_{\{\varphi \leq H_P - s\}} (dd^c \varphi)^d.$$

Indeed, the comparison principle ([11, Corollary 3.6]) and the inclusions of sets above give

$$\int_{\{u \leq H_P - 2s\}} (dd^c v)^d \leq \int_{\{\varphi \leq \frac{v + H_P}{2} - s\}} (dd^c v)^d \leq 2^d \int_{\{\varphi \leq \frac{v + H_P}{2} - s\}} \left( dd^c \frac{v + H_P}{2} \right)^d \\ \leq 2^d \int_{\{\varphi \leq \frac{v + H_P}{2} - s\}} (dd^c \varphi)^d \leq 2^d \int_{\{\varphi \leq H_P - s\}} (dd^c \varphi)^d.$$

The claim is proved. Using (3.3) and (3.2) we obtain

$$\int_{\{u \leq H_P - 2t\}} (H_P - u)(dd^c v)^d \\ \leq 2^{d+1} t \int_{\{\varphi \leq H_P - t\}} (dd^c \varphi)^d + 2^{d+1} \int_t^{+\infty} \left( \int_{\{\varphi \leq H_P - s\}} (dd^c \varphi)^d \right) ds \\ = 2^{d+1} \int_{\{\varphi \leq H_P - t\}} (H_P - \varphi)(dd^c \varphi)^d.$$

□

**Proposition 3.3.** *Let  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$  with  $(dd^c u)^d = \mu$  having support in a nonpluripolar compact set  $K$  so that  $\int_K u d\mu > -\infty$  from Proposition 2.11. Let  $\{Q_j\}$  be a sequence of continuous functions on  $K$  decreasing to  $u$  on  $K$ . Then  $u_j := V_{P,K,Q_j}^* \downarrow u$  on  $\mathbb{C}^d$  and  $\mu_j := (dd^c u_j)^d$  is supported in  $K$ . In particular,  $\mu_j \rightarrow \mu = (dd^c u)^d$  weak-\*. Moreover,*

$$(3.4) \quad \lim_{j \rightarrow \infty} \int_K Q_j d\mu_j = \lim_{j \rightarrow \infty} \int_K Q_j d\mu = \int_K u d\mu > -\infty.$$

*Proof.* We can assume  $\{Q_j\}$  are defined and decreasing to  $u$  on the closure of a bounded open neighborhood  $\Omega$  of  $K$ . By adding a negative constant we can assume that  $Q_1 \leq 0$  on  $\Omega$ . Since  $\{Q_j\}$  is decreasing, so is the sequence  $\{u_j\}$ . Moreover, by [4, Proposition 5.1]  $u_j \leq Q_j$  on  $K \setminus E_j$  where  $E_j$  is pluripolar. But  $u$  is a competitor in the definition of  $V_{P,K,Q_j}^*$  so that  $u \leq u_j$  on  $\mathbb{C}^d$ . Thus  $\tilde{u} := \lim_{j \rightarrow \infty} u_j \geq u$  everywhere and  $\tilde{u} \leq u$  on  $K \setminus E$ , where  $E := \cup_j E_j$  is a pluripolar set. Since  $(dd^c u)^d$  put no mass on pluripolar sets,

$$\int_{\{u < \tilde{u}\}} (dd^c u)^d \leq \int_{E \cup (\mathbb{C}^d \setminus K)} (dd^c u)^d = 0.$$

It thus follows from Proposition 3.1 that  $\tilde{u} \leq u$ , hence  $\tilde{u} = u$  on  $\mathbb{C}^d$ .

The second equality in (3.4) follows from the monotone convergence theorem. It remains to prove that

$$\lim_{j \rightarrow \infty} \int_K (-Q_j) d\mu_j = \int_K (-u) d\mu.$$

For each  $k$  fixed and  $j \geq k$  we have

$$\int_K (-Q_j) d\mu_j \geq \int_K (-Q_k) d\mu_j = \int_{\Omega} (-Q_k) d\mu_j,$$

hence  $\liminf_{j \rightarrow \infty} \int_K (-Q_j) d\mu_j \geq \int_K (-Q_k) d\mu$  since  $\Omega$  is open and  $\mu_j, \mu$  are supported on  $K$ . Letting  $k \rightarrow +\infty$  we arrive at

$$\liminf_{j \rightarrow \infty} \int_K (-Q_j) d\mu_j \geq \int_K (-u) d\mu.$$

It remains to prove that

$$\limsup_{j \rightarrow \infty} \int_K (-Q_j) d\mu_j \leq \int_K (-u) d\mu.$$

The sequence  $\{u_j\}$  is not necessarily uniformly bounded below on  $K$ . However, using the facts that  $Q_j \geq u$  and  $H_P$  is continuous in  $\mathbb{C}^d$ , it



suffices to prove that

$$(3.5) \quad \limsup_{j \rightarrow \infty} \int_K (H_P - u)(dd^c u_j)^d \leq \int_K (H_P - u)(dd^c u)^d.$$

To verify (3.5), we use Lemma 3.2.

By adding a negative constant we can assume that  $u_j \leq H_P$ . For a function  $v$  and for  $t > 0$  we define  $v^t := \max(v, H_P - t)$ . Note that for each  $t$  the sequence  $\{u_j^t\}$  is locally uniformly bounded below. Define

$$a(t) := 2^{d+1} \int_{\{u \leq H_P - t/2\}} (H_P - u)(dd^c u)^d.$$

Since  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$ , from Proposition 2.11 we have  $a(t) \rightarrow 0$  as  $t \rightarrow +\infty$ . By Lemma 3.2 we have

$$(3.6) \quad \sup_{j \geq 1} \int_{\{u \leq H_P - t\}} (H_P - u)(dd^c u_j)^d \leq a(t).$$

By the plurifine property of non-pluripolar Monge-Ampère measures [10, Proposition 1.4] and (3.6) we have

$$\begin{aligned} \int_K (H_P - u)(dd^c u_j)^d &\leq \int_{K \cap \{u > H_P - t\}} (H_P - u)(dd^c u_j)^d + a(t) \\ &= \int_{K \cap \{u > H_P - t\}} (H_P - u^t)(dd^c u_j^t)^d + a(t) \\ &\leq \int_K (H_P - u^t)(dd^c u_j^t)^d + a(t). \end{aligned}$$

Since  $H_P$  is bounded in  $\Omega$ , it follows from [16, Theorem 4.26] that the sequence of positive Radon measures  $(H_P - u^t)(dd^c u_j^t)^d$  converges weakly on  $\Omega$  to  $(H_P - u^t)(dd^c u^t)^d$ . Since  $K$  is compact it then follows that

$$\limsup_j \int_K (H_P - u)(dd^c u_j)^d \leq \int_K (H_P - u^t)(dd^c u^t)^d + a(t).$$

We finally let  $t \rightarrow +\infty$  to conclude the proof in the following manner:

$$\begin{aligned} \int_K (H_P - u^t)(dd^c u^t)^d &\leq \int_{K \cap \{u > H_P - t\}} (H_P - u^t)(dd^c u^t)^d + a(t) \\ &\leq \int_K (H_P - u)(dd^c u)^d + a(t), \end{aligned}$$

where in the first estimate we have used  $\{u \leq H_P - t\} = \{u^t \leq H_P - t\}$  and Lemma 3.2 and in the last estimate we use again the plurifine property.  $\square$

We now give an alternate description of the Legendre-type transform  $E^*$  from (2.9) which will be related to the the rate function in a large deviation principle. Given  $K \subset \mathbb{C}^d$  compact, we let  $\mathcal{M}_P(K)$  denote the space of positive measures on  $K$  of total mass  $\gamma_d$  and we let  $C(K)$  denote the set of continuous, real-valued functions on  $K$ .

**Proposition 3.4.** *Let  $K$  be a nonpluripolar compact set and  $\mu \in \mathcal{M}_P(K)$ . Then*

$$E^*(\mu) = \sup_{v \in C(K)} [E(V_{P,K,v}^*) - \int_K v d\mu].$$

*Proof.* We first treat the case when  $E^*(\mu) = +\infty$ . By Theorem 2.13 there exists  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$  such that  $\int_K u d\mu = -\infty$ . We take a decreasing sequence  $Q_j \in C(K)$  such that  $Q_j \downarrow u$  on  $K$  and set  $u_j := V_{P,K,Q_j}^*$ . Then  $\{u_j\}$  are decreasing; since  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$  and  $E$  is non-decreasing,  $\{E(u_j)\}$  is uniformly bounded and we obtain

$$E(V_{P,K,Q_j}^*) - \int_K Q_j d\mu \rightarrow +\infty,$$

proving the proposition in this case.

Assume now that  $E^*(\mu) < +\infty$ . Theorem 2.13 ensures that  $\int_{\mathbb{C}^d} u d\mu > -\infty$  for all  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$ . By Lemma 2.15,  $\mu$  puts no mass on pluripolar sets. From monotonicity of  $E$  and the definition of  $E^*$  in (2.9) we have

$$E^*(\mu) \geq \sup_{v \in C(K)} [E(V_{P,K,v}^*) - \int_K v d\mu].$$

Here we have used that

$$V_{P,K,v}^* \leq v \text{ q.e. on } K \text{ for } v \in C(K).$$

For the reverse inequality, fix  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$ . Let  $\{Q_j\}$  be a sequence of continuous functions on  $K$  decreasing to  $u$  on  $K$  and set  $u_j := V_{P,K,Q_j}^*$ . Given  $\epsilon > 0$ , we can choose  $j$  sufficiently large so that, by monotone convergence,

$$\int_K Q_j d\mu \leq \int_K u d\mu + \epsilon;$$

and, by monotonicity of  $E$ ,

$$E(V_{P,K,Q_j}^*) \geq E(u).$$

Hence

$$E(V_{P,K,Q_j}^*) - \int_K Q_j d\mu \geq E(u) - \int_K u d\mu - \epsilon$$

so that

$$\sup_{v \in C(K)} [E(V_{P,K,v}^*) - \int_K v d\mu] \geq E^*(\mu)$$

and equality holds. □

**3.2. Transfinite diameter.** Let  $d_n = d_n(P)$  denote the dimension of the vector space  $Poly(nP)$ . We write

$$Poly(nP) = \text{span}\{e_1, \dots, e_{d_n}\}$$

where  $\{e_j(z) := z^{\alpha(j)}\}_{j=1, \dots, d_n}$  are the standard basis monomials. Given  $\zeta_1, \dots, \zeta_{d_n} \in \mathbb{C}^d$ , let

$$(3.7) \quad \begin{aligned} VDM(\zeta_1, \dots, \zeta_{d_n}) &:= \det[e_i(\zeta_j)]_{i,j=1, \dots, d_n} \\ &= \det \begin{bmatrix} e_1(\zeta_1) & e_1(\zeta_2) & \dots & e_1(\zeta_{d_n}) \\ \vdots & \vdots & \ddots & \vdots \\ e_{d_n}(\zeta_1) & e_{d_n}(\zeta_2) & \dots & e_{d_n}(\zeta_{d_n}) \end{bmatrix} \end{aligned}$$

and for  $K \subset \mathbb{C}^d$  compact let

$$V_n = V_n(K) := \max_{\zeta_1, \dots, \zeta_{d_n} \in K} |VDM(\zeta_1, \dots, \zeta_{d_n})|.$$

It was shown in [2] that

$$(3.8) \quad \delta(K) := \delta(K, P) := \lim_{n \rightarrow \infty} V_n^{1/l_n}$$

exists where

$$l_n := \sum_{j=1}^{d_n} \deg(e_j) = \sum_{j=1}^{d_n} |\alpha(j)|$$

is the sum of the degrees of the basis monomials for  $Poly(nP)$ . We call  $\delta(K)$  the  $P$ -transfinite diameter of  $K$ . More generally, for  $w$  an admissible weight function on  $K$  and  $\zeta_1, \dots, \zeta_{d_n} \in K$ , let

$$(3.9) \quad VDM_n^Q(\zeta_1, \dots, \zeta_{d_n}) := VDM(\zeta_1, \dots, \zeta_{d_n}) w(\zeta_1)^n \cdots w(\zeta_{d_n})^n$$

$$= \det \begin{bmatrix} e_1(\zeta_1) & e_1(\zeta_2) & \cdots & e_1(\zeta_{d_n}) \\ \vdots & \vdots & \ddots & \vdots \\ e_{d_n}(\zeta_1) & e_{d_n}(\zeta_2) & \cdots & e_{d_n}(\zeta_{d_n}) \end{bmatrix} \cdot w(\zeta_1)^n \cdots w(\zeta_{d_n})^n$$

be a *weighted Vandermonde determinant*. Let

$$W_n(K) := \max_{\zeta_1, \dots, \zeta_{d_n} \in K} |VDM_n^Q(\zeta_1, \dots, \zeta_{d_n})|.$$

An  $n$ -th *weighted P-Fekete set* for  $K$  and  $w$  is a set of  $d_n$  points  $\zeta_1, \dots, \zeta_{d_n} \in K$  with the property that

$$|VDM_n^Q(\zeta_1, \dots, \zeta_{d_n})| = W_n(K).$$

The limit

$$\delta^Q(K) := \delta^Q(K, P) := \lim_{n \rightarrow \infty} W_n(K)^{1/l_n}$$

exists and is called the *weighted P-transfinite diameter*. The following was proved in [2].

**Theorem 3.5. [Asymptotic Weighted P-Fekete Measures]** *Let  $K \subset \mathbb{C}^d$  be compact with admissible weight  $w$ . For each  $n$ , take points  $z_1^{(n)}, z_2^{(n)}, \dots, z_{d_n}^{(n)} \in K$  for which*

$$(3.10) \quad \lim_{n \rightarrow \infty} [|VDM_n^Q(z_1^{(n)}, \dots, z_{d_n}^{(n)})|]^{1/l_n} = \delta^Q(K)$$

(asymptotically weighted P-Fekete arrays) and let  $\mu_n := \frac{1}{d_n} \sum_{j=1}^{d_n} \delta_{z_j^{(n)}}$ .

Then

$$\mu_n \rightarrow \frac{1}{\gamma_d} \mu_{K,Q} \text{ weak-}^*.$$

Another ingredient we will use is a Rumely-type relation between transfinite diameter and energy of  $V_{P,K,Q}^*$  from [2].

**Theorem 3.6.** *Let  $K \subset \mathbb{C}^d$  be compact and  $w = e^{-Q}$  with  $Q \in C(K)$ . Then*

$$(3.11) \quad \log \delta^Q(K) = \frac{-1}{\gamma_d d A} \mathcal{E}(V_{P,K,Q}^*, H_P) = \frac{-(d+1)}{\gamma_d d A} E(V_{P,K,Q}^*).$$

Here  $A = A(P, d)$  was defined in [2]; we recall the definition. For  $P = \Sigma$  so that  $\text{Poly}(n\Sigma) = \mathcal{P}_n$ , we have

$$d_n(\Sigma) = \binom{d+n}{d} = O(n^d/d!) \text{ and } l_n(\Sigma) = \frac{d}{d+1} n d_n(\Sigma).$$

For a convex body  $P \subset (\mathbb{R}^+)^d$ , define  $f_n(d)$  by writing

$$l_n = f_n(d) \frac{nd}{d+1} d_n = f_n(d) \frac{l_n(\Sigma)}{d_n(\Sigma)} d_n.$$

Then the ratio  $l_n/d_n$  divided by  $l_n(\Sigma)/d_n(\Sigma)$  has a limit; i.e.,

$$(3.12) \quad \lim_{n \rightarrow \infty} f_n(d) =: A = A(P, d).$$

**3.3. Bernstein-Markov.** For  $K \subset \mathbb{C}^d$  compact,  $w = e^{-Q}$  an admissible weight function on  $K$ , and  $\nu$  a finite measure on  $K$ , we say that the triple  $(K, \nu, Q)$  satisfies a weighted Bernstein-Markov property if for all  $p_n \in \mathcal{P}_n$ ,

$$(3.13) \quad \|w^n p_n\|_K \leq M_n \|w^n p_n\|_{L^2(\nu)} \text{ with } \limsup_{n \rightarrow \infty} M_n^{1/n} = 1.$$

Here,  $\|w^n p_n\|_K := \sup_{z \in K} |w(z)^n p_n(z)|$  and

$$\|w^n p_n\|_{L^2(\nu)}^2 := \int_K |p_n(z)|^2 w(z)^{2n} d\nu(z).$$

Following [1], given  $P \subset (\mathbb{R}^+)^d$  a convex body, we say that a finite measure  $\nu$  with support in a compact set  $K$  is a Bernstein-Markov measure for the triple  $(P, K, Q)$  if (3.13) holds for all  $p_n \in \text{Poly}(nP)$ .

For any  $P$  there exists  $A = A(P) > 0$  with  $\text{Poly}(nP) \subset \mathcal{P}_{An}$  for all  $n$ . Thus if  $(K, \nu, Q)$  satisfies a weighted Bernstein-Markov property, then  $\nu$  is a Bernstein-Markov measure for  $(P, K, \tilde{Q})$  where  $\tilde{Q} = AQ$ . In particular, if  $\nu$  is a *strong Bernstein-Markov measure* for  $K$ ; i.e., if  $\nu$  is a weighted Bernstein-Markov measure for any  $Q \in C(K)$ , then for any such  $Q$ ,  $\nu$  is a Bernstein-Markov measure for the triple  $(P, K, Q)$ . Strong Bernstein-Markov measures exist for any nonpluripolar compact set; cf., Corollary 3.8 of [9]. The paragraph following this corollary gives a sufficient mass-density type condition for a measure to be a strong Bernstein-Markov measure.

Given  $P$ , for  $\nu$  a finite measure on  $K$  and  $Q \in \mathcal{A}(K)$ , define

$$(3.14) \quad Z_n := Z_n(P, K, Q, \nu) := \int_K \cdots \int_K |VDM_n^Q(z_1, \dots, z_{d_n})|^2 d\nu(z_1) \cdots d\nu(z_{d_n}).$$

The main consequence of using a Bernstein-Markov measure for  $(P, K, Q)$  is the following:

**Proposition 3.7.** *Let  $K \subset \mathbb{C}^d$  be a compact set and let  $Q \in \mathcal{A}(K)$ . If  $\nu$  is a Bernstein-Markov measure for  $(P, K, Q)$  then*

$$(3.15) \quad \lim_{k \rightarrow \infty} Z_n^{\frac{1}{2l_n}} = \delta^Q(K).$$

*Proof.* That  $\limsup_{k \rightarrow \infty} Z_n^{\frac{1}{2l_n}} \leq \delta^Q(K)$  is clear. Observing from (3.7) and (3.9) that, fixing all variables but  $z_j$ ,

$$z_j \rightarrow VDM_n^Q(z_1, \dots, z_j, \dots, z_{d_n}) = w(z_j)^n p_n(z_j)$$

for some  $p_n \in \text{Poly}(nP)$ , to show  $\liminf_{k \rightarrow \infty} Z_n^{\frac{1}{2l_n}} \geq \delta^Q(K)$  one starts with an  $n$ -th weighted  $P$ -Fekete set for  $K$  and  $w$  and repeatedly applies the weighted Bernstein-Markov property.  $\square$

Recall  $\mathcal{M}_P(K)$  is the space of positive measures on  $K$  with total mass  $\gamma_d$ . With the weak-\* topology, this is a separable, complete metrizable space. A neighborhood basis of  $\mu \in \mathcal{M}_P(K)$  can be given by sets

$$(3.16) \quad G(\mu, k, \epsilon) := \left\{ \sigma \in \mathcal{M}_P(K) : \left| \int_K (\text{Re}z)^\alpha (\text{Im}z)^\beta (d\mu - d\sigma) \right| < \epsilon \right. \\ \left. \text{for } 0 \leq |\alpha| + |\beta| \leq k \right\}$$

where  $\text{Re}z = (\text{Re}z_1, \dots, \text{Re}z_n)$  and  $\text{Im}z = (\text{Im}z_1, \dots, \text{Im}z_n)$ .

Given  $\nu$  as in Proposition 3.7, we define a probability measure  $\text{Prob}_n$  on  $K^{d_n}$  via, for a Borel set  $A \subset K^{d_n}$ ,

$$(3.17) \quad \text{Prob}_n(A) := \frac{1}{Z_n} \cdot \int_A |VDM_n^Q(z_1, \dots, z_{d_n})|^2 \cdot d\nu(z_1) \cdots d\nu(z_{d_n}).$$

We immediately obtain the following:

**Corollary 3.8.** *Let  $\nu$  be a Bernstein-Markov measure for  $(P, K, Q)$ . Given  $\eta > 0$ , define*

$$(3.18) \quad A_{n,\eta} := \{(z_1, \dots, z_{d_n}) \in K^{d_n} : |VDM_n^Q(z_1, \dots, z_{d_n})|^2 \geq (\delta^Q(K) - \eta)^{2l_n}\}.$$

*Then there exists  $n^* = n^*(\eta)$  such that for all  $n > n^*$ ,*

$$\text{Prob}_n(K^{d_n} \setminus A_{n,\eta}) \leq \left( 1 - \frac{\eta}{2\delta^Q(K)} \right)^{2l_n}.$$

**Remark 3.9.** Corollary 3.8 was proved in [9], Corollary 3.2, for  $\nu$  a probability measure but an obvious modification works for  $\nu(K) < \infty$ .

Using (3.17), we get an induced probability measure  $\mathbf{P}$  on the infinite product space of arrays  $\chi := \{X = \{x_j^{(n)}\}_{n=1,2,\dots; j=1,\dots,d_n} : x_j^{(n)} \in K\}$ :

$$(\chi, \mathbf{P}) := \prod_{n=1}^{\infty} (K^{d_n}, Prob_n).$$

**Corollary 3.10.** *Let  $\nu$  be a Bernstein-Markov measure for  $(P, K, Q)$ . For  $\mathbf{P}$ -a.e. array  $X = \{x_j^{(n)}\} \in \chi$ ,*

$$\nu_n := \frac{1}{d_n} \sum_{j=1}^{d_n} \delta_{x_j^{(n)}} \rightarrow \frac{1}{\gamma_d} \mu_{K,Q} \text{ weak-}^*.$$

*Proof.* From Theorem 3.5 it suffices to verify for  $\mathbf{P}$ -a.e. array  $X = \{x_j^{(n)}\}$

$$(3.19) \quad \liminf_{n \rightarrow \infty} (|VDM_n^Q(x_1^{(n)}, \dots, x_{d_n}^{(n)})|)^{\frac{1}{l_n}} = \delta^Q(K).$$

Given  $\eta > 0$ , the condition that for a given array  $X = \{x_j^{(n)}\}$  we have

$$\liminf_{n \rightarrow \infty} (|VDM_n^Q(x_1^{(n)}, \dots, x_{d_n}^{(n)})|)^{\frac{1}{l_n}} \leq \delta^Q(K) - \eta$$

means that  $(x_1^{(n)}, \dots, x_{d_n}^{(n)}) \in K^{d_n} \setminus A_{n,\eta}$  for infinitely many  $n$ . Setting

$$E_n := \{X \in \chi : (x_1^{(n)}, \dots, x_{d_n}^{(n)}) \in K^{d_n} \setminus A_{n,\eta}\},$$

we have

$$\mathbf{P}(E_n) \leq Prob_n(K^{d_n} \setminus A_{n,\eta}) \leq (1 - \frac{\eta}{2\delta^Q(K)})^{2l_n}$$

and  $\sum_{n=1}^{\infty} \mathbf{P}(E_n) < +\infty$ . By the Borel-Cantelli lemma,

$$\mathbf{P}(\limsup_{n \rightarrow \infty} E_n) = \mathbf{P}(\bigcap_{n=1}^{\infty} \bigcup_{k \geq n} E_k) = 0.$$

Thus, with probability one, only finitely many  $E_n$  occur, and (3.19) follows.  $\square$

The main goal in the rest of the paper is to verify a stronger probabilistic result – a large deviation principle – and to explain this result in  $P$ -pluripotential-theoretic terms.

4. RELATION BETWEEN  $E^*$  AND  $J, J^Q$  FUNCTIONALS.

We define some functionals on  $\mathcal{M}_P(K)$  using  $L^2$ -type notions which act as a replacement for an energy functional on measures. Then we show these functionals  $\bar{J}(\mu)$  and  $\underline{J}(\mu)$  defined using a “lim sup” and a “lim inf” coincide (see Definitions 4.1 and 4.2); this is the essence of our first proof of the large deviation principle, Theorem 5.1. Using Proposition 3.4, we relate this functional with  $E^*$  from (2.9).

Fix a nonpluripolar compact set  $K$  and a strong Bernstein-Markov measure  $\nu$  on  $K$ . For simplicity, we normalize so that  $\nu$  is a probability measure. Recall then for any  $Q \in C(K)$ ,  $\nu$  is a Bernstein-Markov measure for the triple  $(P, K, Q)$ . Given  $G \subset \mathcal{M}_P(K)$  open, for each  $s = 1, 2, \dots$  we set

$$(4.1) \quad \tilde{G}_s := \{\mathbf{a} = (a_1, \dots, a_s) \in K^s : \frac{\gamma_d}{s} \sum_{j=1}^s \delta_{a_j} \in G\}.$$

Define, for  $n = 1, 2, \dots$ ,

$$J_n(G) := \left[ \int_{\tilde{G}_{d_n}} |VDM_n(\mathbf{a})|^2 d\nu(\mathbf{a}) \right]^{1/2l_n}.$$

**Definition 4.1.** For  $\mu \in \mathcal{M}_P(K)$  we define

$$\bar{J}(\mu) := \inf_{G \ni \mu} \bar{J}(G) \quad \text{where} \quad \bar{J}(G) := \limsup_{n \rightarrow \infty} J_n(G);$$

$$\underline{J}(\mu) := \inf_{G \ni \mu} \underline{J}(G) \quad \text{where} \quad \underline{J}(G) := \liminf_{n \rightarrow \infty} J_n(G).$$

The infima are taken over all neighborhoods  $G$  of the measure  $\mu$  in  $\mathcal{M}_P(K)$ . A priori,  $\bar{J}, \underline{J}$  depend on  $\nu$ . These functionals are nonnegative but can take the value zero. Intuitively, we are taking a “limit” of  $L^2(\nu)$  averages of discrete, equally weighted approximants  $\frac{\gamma_d}{s} \sum_{j=1}^s \delta_{a_j}$  of  $\mu$ . An “ $L^\infty$ ” version of  $\bar{J}, \underline{J}$  was introduced in [8] where  $J_n(G)$  is replaced by

$$(4.2) \quad W_n(G) := \sup_{\mathbf{a} \in \tilde{G}_{d_n}} |VDM_n(\mathbf{a})|^{1/l_n} \geq J_n(G).$$

The weighted versions of these functionals are defined for  $Q \in \mathcal{A}(K)$  using

$$(4.3) \quad J_n^Q(G) := \left[ \int_{\tilde{G}_{d_n}} |VDM_n^Q(\mathbf{a})|^2 d\nu(\mathbf{a}) \right]^{1/2l_n}.$$



**Definition 4.2.** For  $\mu \in \mathcal{M}_P(K)$  we define

$$\begin{aligned}\bar{J}^Q(\mu) &:= \inf_{G \ni \mu} \bar{J}^Q(G) \text{ where } \bar{J}^Q(G) := \limsup_{n \rightarrow \infty} J_n^Q(G); \\ \underline{J}^Q(\mu) &:= \inf_{G \ni \mu} \underline{J}^Q(G) \text{ where } \underline{J}^Q(G) := \liminf_{n \rightarrow \infty} J_n^Q(G).\end{aligned}$$

The uppersemicontinuity of  $\bar{J}$ ,  $\bar{J}^Q$ ,  $\underline{J}$  and  $\underline{J}^Q$  on  $\mathcal{M}_P(K)$  (with the weak-\* topology) follows as in Lemma 3.1 of [8]. Set

$$b_d = b_d(P) := \frac{d+1}{Ad\gamma_d}.$$

**Proposition 4.3.** Fix  $Q \in C(K)$ . Then

- (1)  $\bar{J}^Q(\mu) \leq \delta^Q(K)$ ;
- (2)  $\bar{J}(\mu) = \bar{J}^Q(\mu) \cdot (e^{\int_K Q d\mu})^{b_d}$ ;
- (3)  $\log \bar{J}(\mu) \leq \inf_{v \in C(K)} [\log \delta^v(K) + b_d \int_K v d\mu]$ ;
- (4)  $\log \bar{J}^Q(\mu) \leq \inf_{v \in C(K)} [\log \delta^v(K) + b_d \int_K v d\mu] - b_d \int_K Q d\mu$ .

Properties (1)-(4) also hold for the functionals  $\underline{J}$ ,  $\underline{J}^Q$ .

*Proof.* Property (1) follows from

$$J_n^Q(G) \leq \sup_{\mathbf{a} \in \tilde{G}_{d_n}} |VDM_n^Q(\mathbf{a})|^{1/l_n} \leq \sup_{\mathbf{a} \in K^{d_n}} |VDM_n^Q(\mathbf{a})|^{1/l_n}.$$

The proofs of Corollary 3.4, Proposition 3.5 and Proposition 3.6 of [8] work mutatis mutandis to verify (2), (3) and (4). The relevant estimation, replacing the corresponding one which is two lines above equation (3.2) in [8], is, given  $\epsilon > 0$ , for  $\mathbf{a} \in \tilde{G}_{d_n}$ ,

$$\begin{aligned}(4.4) \quad |VDM_n^Q(\mathbf{a})| e^{\frac{nd_n}{\gamma_d}(-\epsilon - \int_K Q d\mu)} &\leq |VDM_n(\mathbf{a})| \\ &\leq |VDM_n^Q(\mathbf{a})| e^{\frac{nd_n}{\gamma_d}(\epsilon + \int_K Q d\mu)}.\end{aligned}$$

To see this, we first recall that

$$|VDM_n(\mathbf{a})| = |VDM_n^Q(\mathbf{a})| e^{n \sum_{j=1}^{d_n} Q(a_j)}.$$

For  $\mu \in \mathcal{M}_P(K)$ ,  $Q \in C(K)$ ,  $\epsilon > 0$ , there exists a neighborhood  $G$  of  $\mu$  in  $\mathcal{M}_P(K)$  with

$$-\epsilon < \int_K Q d\mu - \frac{\gamma_d}{d_n} \sum_{j=1}^{d_n} Q(a_j) < \epsilon$$

for  $\mathbf{a} \in \tilde{G}_{d_n}$ . Plugging this double inequality into the previous equality we get (4.4). Moreover, from (3.12),

$$(4.5) \quad \lim_{n \rightarrow \infty} \frac{nd_n}{l_n} = \frac{d+1}{Ad} = b_d \gamma_d$$

so that  $\frac{nd_n}{\gamma_d} \asymp l_n b_d$  as  $n \rightarrow \infty$ . Taking  $l_n$ —the roots in (4.4) accounts for the factor of  $b_d$  in (2), (3) and (4).  $\square$

**Remark 4.4.** The corresponding  $\underline{W}, \underline{W}^Q, \overline{W}, \overline{W}^Q$  functionals, defined using (4.2), clearly dominate their “ $J$ ” counterparts; e.g.,  $\overline{W}^Q \geq \overline{J}^Q$ .

Note that formula (3.11) can be rewritten:

$$(4.6) \quad \log \delta^Q(K) = -b_d E(V_{P,K,Q}^*).$$

Thus the upper bound in Proposition 4.3 (3) becomes

$$(4.7) \quad \log \overline{J}(\mu) \leq -b_d \sup_{v \in C(K)} [E(V_{P,K,v}^*) - \int_K v d\mu] = -b_d E^*(\mu).$$

For the rest of section 4 and section 5, we will always assume  $Q \in C(K)$ . Theorem 4.5 shows that the inequalities in (3) and (4) are equalities, and that the  $\overline{J}, \overline{J}^Q$  functionals coincide with their  $\underline{J}, \underline{J}^Q$  counterparts. The key step in the proof of Theorem 4.5 is to verify this for  $\overline{J}^v(\mu_{K,v})$  and  $\underline{J}^v(\mu_{K,v})$ .

**Theorem 4.5.** *Let  $K \subset \mathbb{C}^d$  be a nonpluripolar compact set and let  $v$  satisfy a strong Bernstein-Markov property. Fix  $Q \in C(K)$ . Then for any  $\mu \in \mathcal{M}_P(K)$ ,*

$$(4.8) \quad \log \overline{J}(\mu) = \log \underline{J}(\mu) = \inf_{v \in C(K)} [\log \delta^v(K) + b_d \int_K v d\mu]$$

and

$$(4.9) \quad \log \overline{J}^Q(\mu) = \log \underline{J}^Q(\mu) = \inf_{v \in C(K)} [\log \delta^v(K) + b_d \int_K v d\mu] - b_d \int_K Q d\mu.$$

*Proof.* It suffices to prove (4.8) since (4.9) follows from (2) of Proposition 4.3. We have the upper bound

$$\log \overline{J}(\mu) \leq \inf_{v \in C(K)} [\log \delta^v(K) + b_d \int_K v d\mu]$$

from (3); for the lower bound, we consider different cases.

Case I:  $\mu = \mu_{K,v}$  for some  $v \in C(K)$ .

We verify that

$$(4.10) \quad \log \bar{J}(\mu_{K,v}) = \log \underline{J}(\mu_{K,v}) = \log \delta^v(K) + b_d \int_K v d\mu_{K,v}$$

which proves (4.8) in this case.

To prove (4.10), we use the definition of  $\underline{J}(\mu_{K,v})$  and Corollary 3.8. Fix a neighborhood  $G$  of  $\mu_{K,v}$ . For  $\eta > 0$ , define  $A_{n,\eta}$  as in (3.18) with  $Q = v$ . Set

$$(4.11) \quad \eta_n := \max \left( \delta^v(K) - \frac{nZ_n^{1/2l_n}}{n+1}, \frac{Z_n^{1/2l_n}}{n+1} \right).$$

By Proposition 3.7,  $\eta_n \rightarrow 0$ . We claim that we have the inclusion

$$(4.12) \quad A_{n,\eta_n} \subset \tilde{G}_{d_n} \text{ for all } n \text{ large enough.}$$

We prove (4.12) by contradiction: if false, there is a sequence  $\{n_j\}$  with  $n_j \uparrow \infty$  and  $x^j = (x_1^j, \dots, x_{d_{n_j}}^j) \in A_{n_j, \eta_{n_j}} \setminus \tilde{G}_{d_{n_j}}$ . However  $\mu_j := \frac{\gamma_d}{d_{n_j}} \sum_{i=1}^{d_{n_j}} \delta_{x_i^j} \notin G$  for  $j$  sufficiently large contradicts Theorem 3.5 since  $x^j \in A_{n_j, \eta_j}$  and  $\eta_j \downarrow 0$  imply  $\mu_j \rightarrow \mu_{K,v}$  weak-\*

Next, a direct computation using (4.11) shows that, for all  $n$  large enough,

$$(4.13) \quad \text{Prob}_n(K^{d_n} \setminus A_{n,\eta_n}) \leq \frac{(\delta^v(K) - \eta_n)^{2l_n}}{Z_n} \leq \left(\frac{n}{n+1}\right)^{2l_n} \leq \frac{n}{n+1}$$

(recall  $\nu$  is a probability measure). Hence

$$\begin{aligned} & \frac{1}{Z_n} \int_{\tilde{G}_{d_n}} |VDM_n^v(z_1, \dots, z_{d_n})|^2 \cdot d\nu(z_1) \cdots d\nu(z_{d_n}) \\ & \geq \frac{1}{Z_n} \int_{A_{n,\eta_n}} |VDM_n^v(z_1, \dots, z_{d_n})|^2 \cdot d\nu(z_1) \cdots d\nu(z_{d_n}) \\ & \geq \frac{1}{n+1}. \end{aligned}$$

Since  $P \subset r\Sigma$  and  $\Sigma \subset kP$  for some  $k \in \mathbb{Z}^+$ ,  $l_n = 0(n^{d+1})$  and we have  $\frac{1}{2l_n} \log(n+1) \rightarrow 0$ . Since  $\nu$  satisfies a strong Bernstein-Markov property and  $v \in C(K)$ , using Proposition 3.7 and the above estimate

we conclude that

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{1}{2l_n} \log \int_{\tilde{G}_{d_n}} |VDM_n^v(z_1, \dots, z_{d_n})|^2 d\nu(z_1) \cdots d\nu(z_{d_n}) \\ \geq \log \delta^v(K). \end{aligned}$$

Taking the infimum over all neighborhoods  $G$  of  $\mu_{K,v}$  we obtain

$$\log \underline{J}^v(\mu_{K,v}) \geq \log \delta^v(K).$$

From (1) Proposition 4.3,  $\log \bar{J}^v(\mu_{K,v}) \leq \log \delta^v(K)$ ; thus we have

$$(4.14) \quad \log \underline{J}^v(\mu_{K,v}) = \log \bar{J}^v(\mu_{K,v}) = \log \delta^v(K).$$

Using (2) of Proposition 4.3 with  $\mu = \mu_{K,v}$  we obtain (4.10).

*Case II:  $\mu \in \mathcal{M}_P(K)$  with the property that  $E^*(\mu) < \infty$ .*

From Theorem 2.13 and Proposition 2.11 there exists  $u \in L_P(\mathbb{C}^d)$  – indeed,  $u \in \mathcal{E}_P^1(\mathbb{C}^d)$  – with  $\mu = (dd^c u)^d$  and  $\int_K u d\mu > -\infty$ . However, since  $u$  is only usc on  $K$ ,  $\mu$  is not necessarily of the form  $\mu_{K,v}$  for some  $v \in C(K)$ . Taking a sequence of continuous functions  $\{Q_j\} \subset C(K)$  with  $Q_j \downarrow u$  on  $K$ , by Proposition 3.3 the weighted extremal functions  $V_{P,K,Q_j}^*$  decrease to  $u$  on  $\mathbb{C}^d$ ;

$$\mu_j := (dd^c V_{P,K,Q_j}^*)^d \rightarrow \mu = (dd^c u)^d \text{ weak-*};$$

and

$$(4.15) \quad \lim_{j \rightarrow \infty} \int_K Q_j d\mu_j = \lim_{j \rightarrow \infty} \int_K Q_j d\mu = \int_K u d\mu.$$

From the previous case we have

$$\log \bar{J}(\mu_j) = \log \underline{J}(\mu_j) = \log \delta^{Q_j}(K) + b_d \int_K Q_j d\mu_j.$$

Using uppersemicontinuity of the functional  $\mu \rightarrow \underline{J}(\mu)$ ,

$$\limsup_{j \rightarrow \infty} \bar{J}(\mu_j) = \limsup_{j \rightarrow \infty} \underline{J}(\mu_j) \leq \underline{J}(\mu).$$

Since  $Q_j \downarrow u$  on  $K$ ,

$$(4.16) \quad \limsup_{j \rightarrow \infty} \log \delta^{Q_j}(K) = \lim_{j \rightarrow \infty} \log \delta^{Q_j}(K).$$

Therefore

$$M := \lim_{j \rightarrow \infty} \log \underline{J}(\mu_j) = \lim_{j \rightarrow \infty} (\log \delta^{Q_j}(K) + b_d \int_K Q_j d\mu_j)$$

exists and is less than or equal to  $\log \underline{J}(\mu)$ . We want to show that

$$(4.17) \quad \inf_v [\log \delta^v(K) + b_d \int_K v d\mu] \leq M.$$

Given  $\epsilon > 0$ , by (4.15) for  $j \geq j_0(\epsilon)$ ,

$$\int_K Q_j d\mu_j \geq \int_K Q_j d\mu - \epsilon \text{ and } \log \underline{J}(\mu_j) < M + \epsilon.$$

Hence for such  $j$ ,

$$\begin{aligned} \inf_v [\log \delta^v(K) + b_d \int_K v d\mu] &\leq \log \delta^{Q_j}(K) + b_d \int_K Q_j d\mu \\ &\leq \log \delta^{Q_j}(K) + b_d \int_K Q_j d\mu_j + b_d \epsilon = \log \underline{J}(\mu_j) + b_d \epsilon < M + (b_d + 1)\epsilon, \end{aligned}$$

yielding (4.17). This finishes the proof in Case II.

*Case III:*  $\mu \in \mathcal{M}(K)$  with the property that  $E^*(\mu) = +\infty$ .

It follows from Proposition 3.4 and Theorem 3.6 that the right-hand side of (4.8) is  $-\infty$ , finishing the proof.  $\square$

**Remark 4.6.** From now on, we simply use the notation  $J, J^Q$  without the overline or underline. Using Proposition 3.4 and Theorem 3.6, we have

$$\begin{aligned} \log J(\mu) &= \inf_{Q \in \mathcal{C}(K)} [\log \delta^Q(K) + b_d \int_K Q d\mu] \\ &= - \sup_{Q \in \mathcal{C}(K)} [-\log \delta^Q(K) - b_d \int_K Q d\mu] \\ &= - \sup_{Q \in \mathcal{C}(K)} [b_d E(V_{P,K,Q}^*) - b_d \int_K Q d\mu] = -b_d \sup_{Q \in \mathcal{C}(K)} [E(V_{P,K,Q}^*) - \int_K Q d\mu] \end{aligned}$$

(recall (4.6)) which one can compare with

$$E^*(\mu) = \sup_{Q \in \mathcal{C}(K)} [E(V_{P,K,Q}^*) - \int_K Q d\mu]$$

from Proposition 3.4 to conclude

$$(4.18) \quad \log J(\mu) = -b_d E^*(\mu).$$

In particular,  $J, J^Q$  are independent of the choice of strong Bernstein-Markov measure for  $K$ .

Following the idea in Proposition 4.3 of [9], we observe the following:

**Proposition 4.7.** *Let  $K \subset \mathbb{C}^d$  be a nonpluripolar compact set and let  $\nu$  satisfy a strong Bernstein-Markov property. Fix  $Q \in C(K)$ . The measure  $\mu_{K,Q}$  is the unique maximizer of the functional  $\mu \rightarrow J^Q(\mu)$  over  $\mu \in \mathcal{M}_P(K)$ ; i.e.,*

$$(4.19) \quad J^Q(\mu_{K,Q}) = \delta^Q(K) \text{ (and } J(\mu_K) = \delta(K)\text{)}.$$

*Proof.* The fact that  $\mu_{K,Q}$  maximizes  $J^Q$  (and  $\mu_K$  maximizes  $J$ ) follows from (4.10), (4.14) and Proposition 4.3.

Assume now that  $\mu \in \mathcal{M}_P(K)$  maximizes  $J^Q$ . From Remark 4.4 and the definitions of the functionals, for any neighborhood  $G \subset \mathcal{M}_P(K)$  of  $\mu$ ,

$$\bar{J}^Q(\mu) \leq \bar{W}^Q(\mu) \leq \sup\{\limsup_{n \rightarrow \infty} |VDM_n^Q(\mathbf{a}^{(n)})|^{1/l_n}\} \leq \delta^Q(K)$$

where the supremum is taken over all arrays  $\{\mathbf{a}^{(n)}\}_{n=1,2,\dots}$  of  $d_n$ -tuples  $\mathbf{a}^{(n)}$  in  $K$  whose normalized counting measures  $\mu_n := \frac{1}{d_n} \sum_{j=1}^{d_n} \delta_{a_j^{(n)}}$  lies in  $G$ . Since  $\bar{J}^Q(\mu) = \delta^Q(K)$  there is an asymptotic weighted Fekete array  $\{\mathbf{a}^{(n)}\}$  as in (3.10). Theorem 3.5 yields that  $\mu_n := \frac{1}{d_n} \sum_{j=1}^{d_n} \delta_{a_j^{(n)}}$  converges weak-\* to  $\mu_{K,Q}$ , hence  $\mu_{K,Q} \in \bar{G}$ . Since this is true for each neighborhood  $G \subset \mathcal{M}_P(K)$  of  $\mu$ , we must have  $\mu = \mu_{K,Q}$ .  $\square$

## 5. LARGE DEVIATION.

As in the previous section, we fix  $K \subset \mathbb{C}^d$  a nonpluripolar compact set;  $Q \in C(K)$ ; and a measure  $\nu$  on  $K$  satisfying a strong Bernstein-Markov property. For  $x_1, \dots, x_{d_n} \in K$ , we get a discrete measure  $\frac{\gamma_d}{d_n} \sum_{j=1}^{d_n} \delta_{x_j} \in \mathcal{M}_P(K)$ . Define  $j_n : K^{d_n} \rightarrow \mathcal{M}_P(K)$  via

$$j_n(x_1, \dots, x_{d_n}) := \frac{\gamma_d}{d_n} \sum_{j=1}^{d_n} \delta_{x_j}.$$

From (3.17),  $\sigma_n := (j_n)_*(\text{Prob}_n)$  is a probability measure on  $\mathcal{M}_P(K)$ : for a Borel set  $B \subset \mathcal{M}_P(K)$ ,

$$(5.1) \quad \sigma_n(B) = \frac{1}{Z_n} \int_{\tilde{B}_{d_n}} |VDM_n^Q(x_1, \dots, x_{d_n})|^2 d\nu(x_1) \cdots d\nu(x_{d_n})$$

where  $\tilde{B}_{d_n} := \{\mathbf{a} = (a_1, \dots, a_{d_n}) \in K^{d_n} : \frac{\gamma_d}{d_n} \sum_{j=1}^{d_n} \delta_{a_j} \in B\}$  (recall (4.1)). Here,  $Z_n := Z_n(P, K, Q, \nu)$ . Note that

$$(5.2) \quad \sigma_n(B)^{1/2l_n} = \frac{1}{Z_n^{1/2l_n}} \cdot J_n^Q(B).$$

For future use, suppose we have a function  $F : \mathbb{R} \rightarrow \mathbb{R}$  and a function  $v \in C(K)$ . We write, for  $\mu \in \mathcal{M}_P(K)$ ,

$$\langle v, \mu \rangle := \int_K v d\mu$$

and then

$$(5.3) \quad \int_{\mathcal{M}_P(K)} F(\langle v, \mu \rangle) d\sigma_n(\mu) := \frac{1}{Z_n} \int_K \cdots \int_K |VDM_n^Q(x_1, \dots, x_{d_n})|^2 F\left(\frac{\gamma_d}{d_n} \sum_{j=1}^{d_n} v(x_j)\right) d\nu(x_1) \cdots d\nu(x_{d_n}).$$

With this notation, we offer two proofs of our LDP, Theorem 5.1. We state the result; define LDP in Definition 5.2; and then proceed with the proofs. This closely follows the exposition in section 5 of [9].

**Theorem 5.1.** *The sequence  $\{\sigma_n = (j_n)_*(\text{Prob}_n)\}$  of probability measures on  $\mathcal{M}_P(K)$  satisfies a **large deviation principle** with speed  $2l_n$  and good rate function  $\mathcal{I} := \mathcal{I}_{K,Q}$  where, for  $\mu \in \mathcal{M}_P(K)$ ,*

$$\mathcal{I}(\mu) := \log J^Q(\mu_{K,Q}) - \log J^Q(\mu).$$

This means that  $\mathcal{I} : \mathcal{M}_P(K) \rightarrow [0, \infty]$  is a lowersemicontinuous mapping such that the sublevel sets  $\{\mu \in \mathcal{M}_P(K) : \mathcal{I}(\mu) \leq \alpha\}$  are compact in the weak-\* topology on  $\mathcal{M}_P(K)$  for all  $\alpha \geq 0$  ( $\mathcal{I}$  is “good”) satisfying (5.4) and (5.5):

**Definition 5.2.** The sequence  $\{\mu_k\}$  of probability measures on  $\mathcal{M}_P(K)$  satisfies a **large deviation principle** (LDP) with good rate function  $\mathcal{I}$  and speed  $2l_n$  if for all measurable sets  $\Gamma \subset \mathcal{M}_P(K)$ ,

$$(5.4) \quad - \inf_{\mu \in \Gamma^0} \mathcal{I}(\mu) \leq \liminf_{n \rightarrow \infty} \frac{1}{2l_n} \log \mu_n(\Gamma) \text{ and}$$

$$(5.5) \quad \limsup_{n \rightarrow \infty} \frac{1}{2l_n} \log \mu_n(\Gamma) \leq - \inf_{\mu \in \bar{\Gamma}} \mathcal{I}(\mu).$$

In the setting of  $\mathcal{M}_P(K)$ , to prove a LDP it suffices to work with a base for the weak-\* topology. The following is a special case of a basic general existence result for a LDP given in Theorem 4.1.11 in [13].

**Proposition 5.3.** *Let  $\{\sigma_\epsilon\}$  be a family of probability measures on  $\mathcal{M}_P(K)$ . Let  $\mathcal{B}$  be a base for the topology of  $\mathcal{M}_P(K)$ . For  $\mu \in \mathcal{M}_P(K)$  let*

$$\mathcal{I}(\mu) := - \inf_{\{G \in \mathcal{B}: \mu \in G\}} \left( \liminf_{\epsilon \rightarrow 0} \epsilon \log \sigma_\epsilon(G) \right).$$

Suppose for all  $\mu \in \mathcal{M}_P(K)$ ,

$$\mathcal{I}(\mu) = - \inf_{\{G \in \mathcal{B}: \mu \in G\}} \left( \limsup_{\epsilon \rightarrow 0} \epsilon \log \sigma_\epsilon(G) \right).$$

Then  $\{\sigma_\epsilon\}$  satisfies a LDP with rate function  $\mathcal{I}(\mu)$  and speed  $1/\epsilon$ .

There is a converse to Proposition 5.3, Theorem 4.1.18 in [13]. For  $\mathcal{M}_P(K)$ , it reads as follows:

**Proposition 5.4.** *Let  $\{\sigma_\epsilon\}$  be a family of probability measures on  $\mathcal{M}_P(K)$ . Suppose that  $\{\sigma_\epsilon\}$  satisfies a LDP with rate function  $\mathcal{I}(\mu)$  and speed  $1/\epsilon$ . Then for any base  $\mathcal{B}$  for the topology of  $\mathcal{M}_P(K)$  and any  $\mu \in \mathcal{M}_P(K)$*

$$\begin{aligned} \mathcal{I}(\mu) &:= - \inf_{\{G \in \mathcal{B}: \mu \in G\}} \left( \liminf_{\epsilon \rightarrow 0} \epsilon \log \sigma_\epsilon(G) \right) \\ &= - \inf_{\{G \in \mathcal{B}: \mu \in G\}} \left( \limsup_{\epsilon \rightarrow 0} \epsilon \log \sigma_\epsilon(G) \right). \end{aligned}$$

**Remark 5.5.** Assuming Theorem 5.1, this shows that, starting with a strong Bernstein-Markov measure  $\nu$  and the corresponding sequence of probability measures  $\{\sigma_n\}$  on  $\mathcal{M}_P(K)$  in (5.1), the existence of an LDP with rate function  $\mathcal{I}(\mu)$  and speed  $2l_n$  implies that necessarily

$$(5.6) \quad \mathcal{I}(\mu) = \log J^Q(\mu_{K,Q}) - \log J^Q(\mu).$$

Uniqueness of the rate function is basic (cf., Lemma 4.1.4 of [13]).

We turn to the first proof of Theorem 5.1, using Theorem 4.5, which gives a pluripotential theoretic description of the rate functional.

*Proof.* As a base  $\mathcal{B}$  for the topology of  $\mathcal{M}_P(K)$ , we can take the sets from (3.16) or simply all open sets. For  $\{\sigma_\epsilon\}$ , we take the sequence of probability measures  $\{\sigma_n\}$  on  $\mathcal{M}_P(K)$  and we take  $\epsilon = \frac{1}{2l_n}$ . For  $G \in \mathcal{B}$ , from (5.2),

$$\frac{1}{2l_n} \log \sigma_n(G) = \log J_n^Q(G) - \frac{1}{2l_n} \log Z_n.$$



From Proposition 3.7, and (4.14) with  $v = Q$ ,

$$\lim_{n \rightarrow \infty} \frac{1}{2l_n} \log Z_n = \log \delta^Q(K) = \log J^Q(\mu_{K,Q});$$

and by Theorem 4.5,

$$\inf_{G \ni \mu} \limsup_{n \rightarrow \infty} \log J_n^Q(G) = \inf_{G \ni \mu} \liminf_{n \rightarrow \infty} \log J_n^Q(G) = \log J^Q(\mu).$$

Thus by Proposition 5.3  $\{\sigma_n\}$  satisfies an LDP with rate function

$$\mathcal{I}(\mu) := \log J^Q(\mu_{K,Q}) - \log J^Q(\mu)$$

and speed  $2l_n$ . This rate function is good since  $\mathcal{M}_P(K)$  is compact.  $\square$

**Remark 5.6.** From Proposition 4.7,  $\mu_{K,Q}$  is the unique maximizer of the functional

$$\mu \rightarrow \log J^Q(\mu)$$

over all  $\mu \in \mathcal{M}_P(K)$ . Thus

$$\mathcal{I}_{K,Q}(\mu) \geq 0 \text{ with } \mathcal{I}_{K,Q}(\mu) = 0 \iff \mu = \mu_{K,Q}.$$

To summarize,  $\mathcal{I}_{K,Q}$  is a good rate function with unique minimizer  $\mu_{K,Q}$ . Using the relations

$$\log J(\mu) = -b_d \sup_{Q \in C(K)} [E(V_{P,K,Q}^*) - \int_K Q d\mu]$$

$$J(\mu) = J^Q(\mu) \cdot (e^{\int_K Q d\mu})^{b_d}, \text{ and } J^Q(\mu_{K,Q}) = \delta^Q(K)$$

(the latter from (4.19)), we have

$$\begin{aligned} \mathcal{I}(\mu) &:= \log \delta^Q(K) - \log J^Q(\mu) \\ &= \log \delta^Q(K) - \log J(\mu) + b_d \int_K Q d\mu \\ &= b_d \sup_{Q \in C(K)} [E(V_{P,K,Q}^*) - \int_K Q d\mu] + \log \delta^Q(K) + b_d \int_K Q d\mu \\ &= b_d \sup_{v \in C(K)} [E(V_{P,K,v}^*) - \int_K v d\mu] - b_d [E(V_{P,K,Q}^*) - \int_K Q d\mu] \end{aligned}$$

from (4.6).

The second proof of our LDP follows from Corollary 4.6.14 in [13], which is a general version of the Gärtner-Ellis theorem. This approach was originally brought to our attention by S. Boucksom and was also utilized by R. Berman in [5]. We state the version of the [13] result for an appropriate family of probability measures.

**Proposition 5.7.** *Let  $C(K)^*$  be the topological dual of  $C(K)$ , and let  $\{\sigma_\epsilon\}$  be a family of probability measures on  $\mathcal{M}_P(K) \subset C(K)^*$  (equipped with the weak- $*$  topology). Suppose for each  $\lambda \in C(K)$ , the limit*

$$\Lambda(\lambda) := \lim_{\epsilon \rightarrow 0} \epsilon \log \int_{C(K)^*} e^{\lambda(x)/\epsilon} d\sigma_\epsilon(x)$$

*exists as a finite real number and assume  $\Lambda$  is Gâteaux differentiable; i.e., for each  $\lambda, \theta \in C(K)$ , the function  $f(t) := \Lambda(\lambda + t\theta)$  is differentiable at  $t = 0$ . Then  $\{\sigma_\epsilon\}$  satisfies an LDP in  $C(K)^*$  with the convex, good rate function  $\Lambda^*$ .*

Here

$$\Lambda^*(x) := \sup_{\lambda \in C(K)} (\langle \lambda, x \rangle - \Lambda(\lambda)),$$

is the Legendre transform of  $\Lambda$ . The upper bound (5.5) in the LDP holds with rate function  $\Lambda^*$  under the assumption that the limit  $\Lambda(\lambda)$  exists and is finite; the Gâteaux differentiability of  $\Lambda$  is needed for the lower bound (5.4). To verify this property in our setting, we must recall a result from [2].

**Proposition 5.8.** *For  $Q \in \mathcal{A}(K)$  and  $u \in C(K)$ , let*

$$F(t) := E(V_{P,K,Q+tu}^*)$$

*for  $t \in \mathbb{R}$ . Then  $F$  is differentiable and*

$$F'(t) = \int_{C^d} u(dd^c V_{P,K,Q+tu}^*)^d.$$

In [2] it was assumed that  $u \in C^2(K)$  but the result is true with the weaker assumption  $u \in C(K)$  (cf., Theorem 11.11 in [16] due to Lu and Nguyen [17], see also [11, Proposition 4.20]).

We proceed with the second proof of Theorem 5.1. For simplicity, we normalize so that  $\gamma_d = 1$  to fit the setting of Proposition 5.7 (so members of  $\mathcal{M}_P(K)$  are probability measures).

*Proof.* We show that for each  $v \in C(K)$ ,

$$\Lambda(v) := \lim_{n \rightarrow \infty} \frac{1}{2l_n} \log \int_{C(K)^*} e^{2l_n \langle v, \mu \rangle} d\sigma_n(\mu)$$

exists as a finite real number. First, since  $\sigma_n$  is a measure on  $\mathcal{M}_P(K)$ , the integral can be taken over  $\mathcal{M}_P(K)$ . Consider

$$\frac{1}{2l_n} \log \int_{\mathcal{M}_P(K)} e^{2l_n \langle v, \mu \rangle} d\sigma_n(\mu).$$

By (5.3), this is equal to

$$\frac{1}{2l_n} \log \frac{1}{Z_n} \cdot \int_{K^{d_n}} |VDM_n^{Q - \frac{l_n}{nd_n}v}(x_1, \dots, x_{d_n})|^2 d\nu(x_1) \cdots d\nu(x_{d_n}).$$

From (4.5), with  $\gamma_d = 1$ ,  $\frac{l_n}{nd_n} \rightarrow \frac{1}{b_d}$ ; hence for any  $\epsilon > 0$ ,

$$\frac{1}{b_d + \epsilon} v \leq \frac{l_n}{nd_n} v \leq \frac{1}{b_d - \epsilon} v \text{ on } K$$

for  $n$  sufficiently large. Recall that

$$Z_n = \int_{K^{d_n}} |VDM_n^Q(x_1, \dots, x_{d_n})|^2 d\nu(x_1) \cdots d\nu(x_{d_n}).$$

Define

$$\tilde{Z}_n := \int_{K^{d_n}} |VDM_n^{Q-v/b_d}(x_1, \dots, x_{d_n})|^2 d\nu(x_1) \cdots d\nu(x_{d_n}).$$

Then we have

$$\lim_{n \rightarrow \infty} \tilde{Z}_n^{\frac{1}{2l_n}} = \delta^{Q-v/b_d}(K) \text{ and } \lim_{n \rightarrow \infty} Z_n^{\frac{1}{2l_n}} = \delta^Q(K)$$

from (3.15) in Proposition 3.7 and the assumption that  $(K, \nu, \tilde{Q})$  satisfies the weighted Bernstein-Markov property for *all*  $\tilde{Q} \in C(K)$ . Thus

$$(5.7) \quad \Lambda(v) = \lim_{n \rightarrow \infty} \frac{1}{2l_n} \log \frac{\tilde{Z}_n}{Z_n} = \log \frac{\delta^{Q-v/b_d}(K)}{\delta^Q(K)}.$$

Define now, for  $v, v' \in C(K)$ ,

$$f(t) := E(V_{P,K,Q-(v+tv')}^*).$$

Proposition 5.8 shows that  $\Lambda$  is Gâteaux differentiable and Proposition 5.7 gives that  $\Lambda^*$  is a rate function on  $C(K)^*$ .

Since each  $\sigma_n$  has support in  $\mathcal{M}_P(K)$ , it follows from (5.4) and (5.5) in Definition 5.2 of an LDP with  $\Gamma \subset C(K)^*$  that for  $\mu \in C(K)^* \setminus \mathcal{M}_P(K)$ ,  $\Lambda^*(\mu) = +\infty$ . By Lemma 4.1.5 (b) of [13], the restriction of  $\Lambda^*$  to  $\mathcal{M}_P(K)$  is a rate function. Since  $\mathcal{M}_P(K)$  is compact, it is a good rate function. Being a Legendre transform,  $\Lambda^*$  is convex.

To compute  $\Lambda^*$ , we have, using (5.7) and (3.11),

$$\begin{aligned}\Lambda^*(\mu) &= \sup_{v \in C(K)} \left( \int_K v d\mu - \log \frac{\delta^{Q-v/b_d}(K)}{\delta^Q(K)} \right) \\ &= \sup_{v \in C(K)} \left( \int_K v d\mu - b_d [E(V_{P,K,Q}^*) - E(V_{P,K,Q-v/b_d}^*)] \right).\end{aligned}$$

Thus

$$\begin{aligned}\Lambda^*(\mu) + b_d E(V_{P,K,Q}^*) &= \sup_{v \in C(K)} \left( \int_K v d\mu + b_d E(V_{P,K,Q-v/b_d}^*) \right) \\ &= \sup_{u \in C(K)} \left( b_d E(V_{P,K,Q+u}^*) - b_d \int_K u d\mu \right) \text{ (taking } u = -v/b_d \text{)}.\end{aligned}$$

Rearranging and replacing  $u$  in the supremum by  $v = u + Q$ ,

$$\begin{aligned}\Lambda^*(\mu) &= \sup_{u \in C(K)} \left( b_d E(V_{P,K,Q+u}^*) - b_d \int_K u d\mu \right) - b_d E(V_{P,K,Q}^*) \\ &= b_d \left[ \sup_{v \in C(K)} E(V_{P,K,v}^*) - \int_K v d\mu \right] - b_d \left[ E(V_{P,K,Q}^*) - \int_K Q d\mu \right]\end{aligned}$$

which agrees with the formula in Remark 5.6 (since  $\mu$  is a probability measure).

□

**Remark 5.9.** Thus the rate function can be expressed in several equivalent ways:

$$\begin{aligned}\mathcal{I}(\mu) &= \Lambda^*(\mu) = \log J^Q(\mu_{K,Q}) - \log J^Q(\mu) \\ &= b_d \left[ \sup_{v \in C(K)} E(V_{P,K,v}^*) - \int_K v d\mu \right] - b_d \left[ E(V_{P,K,Q}^*) - \int_K Q d\mu \right] \\ &= b_d E^*(\mu) - b_d \left[ E(V_{P,K,Q}^*) - \int_K Q d\mu \right]\end{aligned}$$

which generalizes the result equating (5.3), (5.10) and (5.11) in [9] for the case  $P = \Sigma$  and  $b_d = 1$ . Note in the last equality we are using the slightly different notion of  $E^*$  in (2.9) and Proposition 3.4 than that used in [9].

## REFERENCES

- [1] T. Bayraktar, Zero distribution of random sparse polynomials, *Mich. Math. J.*, **66**, (2017), no. 2, 389–419.
- [2] T. Bayraktar, T. Bloom, N. Levenberg, Pluripotential theory and convex bodies, *Mat. Sbornik*, **209** (2018), no. 3, 67–101.
- [3] E. Bedford and B. A. Taylor, B. A., The Dirichlet problem for a complex Monge-Ampère equation, *Invent. Math.*, **37**, (1976), no. 1, 1–44.
- [4] E. Bedford and B. A. Taylor, A new capacity for plurisubharmonic functions. *Acta Math.*, **149**, (1982), no. 1-2, 1–40.
- [5] R. Berman, Determinantal Point Processes and Fermions on Complex Manifolds: Large Deviations and Bosonization, *Comm. Math. Phys.*, **327** (2014), no. 1, 1–47.
- [6] R. Berman and S. Boucksom, Growth of balls of holomorphic sections and energy at equilibrium, *Invent. Math.*, **181**, (2010), 337-394.
- [7] R. Berman, S. Boucksom, V. Guedj and A. Zeriahi, A variational approach to complex Monge-Ampère equations, *Publ. Math. de l’IHÉS*, **117**, (2013), 179–245.
- [8] T. Bloom and N. Levenberg, Pluripotential energy, *Potential Analysis*, **36**, no. 1, 155-176, 2012.
- [9] T. Bloom and N. Levenberg, Pluripotential energy and large deviation, *Indiana Univ. Math. J.*, **62**, no. 2, 523-550, 2013.
- [10] S. Boucksom, P. Eyssidieux, V. Guedj and A. Zeriahi, Monge-Ampère equations in big cohomology classes, *Acta Math.*, **205** (2010), 199-262.
- [11] T. Darvas, E. Di Nezza, and C. H. Lu, Monotonicity of nonpluripolar products and complex Monge-Ampère equations with prescribed singularity, *Analysis & PDE*, **11**, (2018), no. 8, 2049-2087.
- [12] T. Darvas, E. Di Nezza, and C. H. Lu, Log-concavity of volume and complex Monge-Ampère equations with prescribed singularity, arXiv:1807.00276.
- [13] A. Dembo and O. Zeitouni, *Large deviations techniques and applications*, Jones and Bartlett Publishers, Boston, MA, 1993.
- [14] V. Guedj and A. Zeriahi, Intrinsic capacities on compact Kähler manifolds, *J. Geom. Anal.*, **15** (2005), no. 4, 607-639.
- [15] V. Guedj and A. Zeriahi, The weighted Monge-Ampère energy of quasiplurisubharmonic functions, *J. Funct. Anal.*, **250** (2007), no. 2, 442-482.
- [16] V. Guedj and A. Zeriahi, *Degenerate Complex Monge-Ampère Equations*, European Math. Soc. Tracts in Mathematics Vol. 26, 2017.
- [17] C.H. Lu, V.D. Nguyen, Degenerate complex Hessian equations on compact Kähler manifolds, *Indiana Univ. Math. J.*, **64** (2015), no. 6, 1721–1745.
- [18] J. Ross and D. Witt Nyström, Analytic test configurations and geodesic rays, *J. Symplectic Geom.*, **12** (2014), no. 1, 125-169.
- [19] E. Saff and V. Totik, *Logarithmic potentials with external fields*, Springer-Verlag, Berlin, 1997.
- [20] David Witt Nyström, Monotonicity of nonpluripolar Monge-Ampère masses, arXiv:1703.01950. To appear in *Indiana University Mathematics Journal*.

SABANCI UNIVERSITY, ISTANBUL, TURKEY

*Email address:* `tbayraktar@sabanciuniv.edu`

UNIVERSITY OF TORONTO, TORONTO, ONTARIO M5S 2E4 CANADA

*Email address:* `bloom@math.toronto.edu`

INDIANA UNIVERSITY, BLOOMINGTON, IN 47405 USA

*Email address:* `nlevenbe@indiana.edu`

UNIVERSITÉ PARIS-SUD, ORSAY, FRANCE, 91405

*Email address:* `hoang-chinh.lu@u-psud.fr`