

Nonlinear interglitch dynamics, the braking index of the Vela pulsar and the time to the next glitch

O. Akbal,¹★ M. A. Alpar,¹ S. Buchner^{2,3,4} and D. Pines⁵

¹Faculty of Engineering and Natural Sciences, Sabanci University, Orhanli, 34956 Istanbul, Turkey

²Square Kilometer Array South Africa, The Park, Park Road, Pinelands, Cape Town 7405, South Africa

³Hartebeesthoek Radio Astronomy Observatory, PO Box 443, Krugersdorp 1740, South Africa

⁴School of Physics, University of the Witwatersrand, PO Box Wits, Johannesburg 2050, South Africa

⁵Santa Fe Institute, 1399 Hyde Park Rd., Santa Fe, NM 87501, USA

Accepted 2017 May 3. Received 2017 May 3; in original form 2016 December 12

ABSTRACT

The interglitch timing of the Vela pulsar is characterized by a constant second derivative of the rotation rate. This takes over after the post-glitch exponential relaxation and is completed at about the time of the next glitch. The vortex creep model explains the second derivatives in terms of nonlinear response to the glitch. We present interglitch timing fits to the present sample covering 16 large glitches, taking into account the possibility that in some glitches part of the step in the spin-down rate may involve a ‘persistent shift’, as observed in the Crab pulsar. Modifying the expression for the time between glitches with this hypothesis leads to better agreement with the observed interglitch time intervals. We extrapolate the interglitch model fits to obtain spin-down rates just prior to each glitch and use these to calculate the braking index $n = 2.81 \pm 0.12$. The next glitch should occur around 2017 December 22, ± 197 d if no persistent shift is involved, but could occur as early as 2016 July 27, ± 152 d if the 2013 glitch gave rise to a typical Vela persistent shift. Note added: Literally while we were submitting the first version of this paper on 2016 December 12, we saw ATel # 9847 announcing a Vela pulsar glitch which has arrived 138 d after our prediction with a persistent shift, within the 1σ uncertainty of 152 d.

Key words: dense matter – stars: interiors – stars: neutron – pulsars: general – pulsars: individuals: Vela – stars: rotation.

1 INTRODUCTION

The Vela pulsar, PSR 0833-45, was the first pulsar for which a glitch, a sudden speed-up, in spin-down behaviour was observed (Radhakrishnan & Manchester 1969; Reichley & Downs 1969), and it continues to be one of the most active glitching pulsars, as 15 additional large glitch events have been observed in the succeeding 47 yr.¹ Because it has been monitored almost daily since 1985, it has proved possible to determine the onset of glitches with an uncertainty of less than a day and to follow in detail its post-glitch behaviour for the 10 glitches that have occurred since that time (McCulloch et al. 1987; Flanagan 1991; Flanagan & McCulloch 1994; Wang et al. 2000; Dodson, McCulloch & Lewis 2002; Dodson et al. 2004; Flanagan & Buchner 2006; Buchner 2010, 2013; Yu et al. 2013). The 2000 glitch, in particular, happened during an observation session, so that an upper limit of 40 s could be placed on the glitch rise time

(Dodson et al. 2002). In 1994, the Vela pulsar exhibited two large glitches with $\Delta\Omega/\Omega = 8.61 \times 10^{-7}$ and $\Delta\Omega/\Omega = 1.99 \times 10^{-7}$ that were separated by just 32 d (Buchner & Flanagan 2011). The glitch rates, signatures and interglitch behaviour of many pulsars, and the statistics of pulsar glitches indicate similarities to the Vela pulsar (Alpar & Baykal 1994, 2006), once one scales with pulsar rotation frequency and spin-down rate. We believe the Vela pulsar is a Rosetta stone for understanding pulsar glitches and the results presented here have general applicability to pulsar dynamics.

It was recognized early on (Baym, Pethick & Pines 1969) that the fact that one is able to see a well-defined glitch relaxation meant the neutrons present in the crust and core of a pulsar must be superfluid. The most successful phenomenological explanation of the origin of pulsar glitches and post-glitch behaviour has been based on the proposals by Packard (1972) and Anderson & Itoh (1975) that glitches are an intrinsic property of the rotating superfluid and originate in the unpinning of vortices that are pinned to crustal nuclei. In this picture, the pinned superfluid will lag the spin-down of the pulsar until a critical angular velocity is reached, at which point the pinned vortices unpin and move rapidly, transferring angular momentum

* E-mail: akbalonur85@gmail.com

¹ <http://www.jb.man.ac.uk/pulsar/glitches/gTable.html>

to the crust, which is observed as the glitch. Apart from the sudden unpinning events, vortices also creep, by thermal activation against the pinning barriers, allowing the superfluid to spin-down (Alpar et al. 1984a,b).

The vortex creep model posits two different kinds of response to a glitch; a linear response characterized by exponential relaxation, and a nonlinear response characterized by a step in the spin-down rate and subsequent interglitch recovery with an approximately constant second derivative of the rotation frequency (Alpar et al. 1984b; Alpar, Cheng & Pines 1989). The model has proven successful in that it provides a natural explanation for the observed transient exponential decays and for the long-term recovery extending until the next glitch, which can only be explained by nonlinear dynamics. The model also provides a rough estimate of the interglitch intervals.

This understanding of the systematic interglitch timing behaviour due to *internal torques* between the crust and superfluid components of the neutron star makes it possible to take account of these internal torques to obtain the braking index n that characterizes the *external* pulsar torque, and, as we shall see, leads to results that are quite different from an analysis that does not take the internal torques fully into account.

Our aim in the present paper is threefold. First, we study the recovery of the glitch $\Delta\dot{\Omega}$ in the spin-down rate. We confirm that after the initial exponential relaxation components, the negative step $\Delta\dot{\Omega}$ recovers with an approximately constant second derivative $\ddot{\Omega}$ of the rotation rate. As observed in the interglitch intervals up to the ninth observed glitch (Alpar et al. 1993) this recovery is completed at about the time of arrival of the next glitch. Indeed, in a multicomponent dynamical system, the response of internal torques to offsets from steady state always leads to eventual recovery of the steady state. The question is whether the next glitch arrives before or after the completion of the recovery after the previous glitch. Motivated by the earlier observation that the recovery was completed at about the time of arrival of the next glitch in the earlier work on nine glitches, we now investigate if the coincidence between the time of completion of the recovery from one glitch and the time of arrival of the next glitch is still observed in the present sample of glitches. This behaviour of complete recovery with constant $\ddot{\Omega}$ is confirmed in all interglitch intervals in the present sample. We confirm, model independently, that the recovery is completed at about the time of arrival of the next glitch. As any power-law behaviour, this recovery reflects some underlying nonlinear dynamics. We next confirm, with the current data set, that the nonlinear creep model continues to fit the data for post-glitch behaviour and enables one to estimate the time intervals between glitches; we find that the accuracy of the estimated times to the next glitch has improved with the doubling of the glitch sample. Secondly, we explore the possibility that some Vela glitches are accompanied by ‘persistent shifts’ in the spin-down rate of the kind observed in the Crab pulsar, where a sudden decrease in the spin-down rate at the time of a glitch is observed to remain constant, with no healing, until the arrival of the next glitch, at which it is superseded with an additional similar shift. This hypothesis leads to somewhat shorter estimates of the time to the next glitch, which agree better with the observed time intervals. Thus, the persistent shift hypothesis improves the accuracy of the estimates. Thirdly, based on our understanding of the internal torques, and in particular the correlation between the time when the nonlinear creep response is completed and the time of the next glitch, supported by our estimates, we note that the best fiducial epoch for determining the braking index n due to the external (pulsar) torque is the epoch when the response of internal torques to the previous glitch have been completed, i.e. just before the next glitch.

Fitting a long-term ‘true’ pulsar second derivative of the rotation rate to the spin-down rates at these epochs, we find that the Vela pulsar’s braking index $n = 2.81 \pm 0.12$, in agreement with most other measured pulsar braking indices, which lie between $n = 1.8$ and $n = 3$ for an isolated ideal dipole rotating in vacuum (Melatos 1997; Antonopoulou et al. 2015; Archibald et al. 2015; Lyne et al. 2015; Clark et al. 2016). In their determination of the Vela pulsar’s braking index, Lyne et al. (1996) assumed the effects of internal torques would be over with the exponential relaxations and took a fiducial time of 150 d after each glitch to derive a much lower ‘anomalous’ braking index, $n = 1.4 \pm 0.2$. Recent work (Espinoza, Lyne & Stappers 2017) takes into account a term with recovery at a constant $\ddot{\Omega}$, but assumes that the recovery is not completed. The braking index, $n = 1.7 \pm 0.2$ is obtained. As we show in this paper the recovery at constant interglitch $\ddot{\Omega}$ is actually completed just before the next glitch. We find $n = 2.81 \pm 0.12$.

The plan of our paper is as follows. Section 2 contains a summary of the vortex creep model that is used in Section 3, to fit all currently available interglitch timing data. Section 4 describes the estimation of interglitch time intervals with or without the inclusion of persistent shifts. Section 5 derives the braking index of the Vela pulsar for several different variants of our model. Section 6 contains our conclusions.

2 THE VORTEX CREEP MODEL

The post-glitch behaviour of the Vela pulsar exhibits both linear response, in the form of three distinct exponential relaxations, and a nonlinear response with constant $\ddot{\Omega}$ that persists until the next glitch. The exponential decays have relaxation times $\tau \lesssim 32$ d. The vortex creep model (Alpar et al. 1984a,b, 1989) explains glitches and post-glitch behaviour in terms of superfluid dynamics that takes into account vortex pinning, unpinning and creep. We summarize the main concepts here, referring the reader to earlier work for details.

The superfluid components of the neutron star rotate by sustaining quantized vortices and their spin-down in response to the pulsar torque is described by the motion of these vortices radially outward from the rotation axis. Vortex motion is impeded by pinning to inhomogeneities, such as the nuclei in the neutron star’s inner crust where the neutron superfluid coexists with the crustal crystalline lattice (Alpar 1977) or to toroidal flux lines in the outer core of the neutron star (Gügercinoğlu & Alpar 2014). Vortices unpin and repin by thermal activation, thereby providing a vortex creep current. Because of pinning, the superfluid rotates somewhat faster than the crust as the crust spins down under the external torque. The lag $\omega \equiv \Omega_s - \Omega_c$ between the rotation rates of the superfluid and the crust provides a bias to drive a vortex creep current in the radially outward direction from the rotation axis. Vortex creep thus allows the superfluid to spin-down. This process has a steady state when both superfluid and normal matter are spinning down at the same rate, driven by a steady state lag ω_∞ .

If the lag reaches the maximum value $\omega_{cr} > \omega_\infty$ that can be sustained by the pinning forces, vortices unpin and move outward rapidly in an avalanche, thereby transferring angular momentum to the crust, leading to the glitch (Packard 1972; Anderson & Itoh 1975). In addition to the parts of the pinned superfluid with continuous vortex current, analogous to resistors in an electric circuit; there are also vortex traps prone to catastrophic unpinning, interspersed with vortex-free regions, analogous to capacitors. Vortex traps are sites of extra pinning strength, where critical conditions for unpinning can be reached due to this enhanced vortex density (Chau

et al. 1993; Mochizuki & Izuyama 1995). The high vortex density in the traps leads to local superfluid velocities that are too large to permit pinned vortices in the regions surrounding the traps, which are therefore vortex-free regions, containing few and ineffective pinning centres. Vortex-free regions contribute to the angular momentum transfer at glitches, but do not contribute to the spin-down between glitches since they do not sustain vortices participating in the creep process.

When critical conditions $\omega = \omega_{\text{cr}}$ are reached, vortices unpin collectively from vortex traps, scatter through the vortex-free regions and incite unpinning at further vortex traps, and so create an avalanche. This sudden transfer of angular momentum to the crust, analogous to charge transfer in capacitor discharges, is observed as a glitch in the steady spin-down of the pulsar. Glitches by critical unpinning superposed on an underlying vortex creep process were simulated by a ‘coherent noise’ statistical model by Melatos & Warszawski (2009), who derive interesting information on pinning parameters by comparison of their simulations with the statistics of frequently glitching pulsars. Further interesting simulations were made by Warszawski & Melatos (2011) in a model where glitches by vortex unpinning are superposed on a continuous relaxation of the superfluid described by the Gross–Pitaevskii Equation. The constraints obtained in these papers on pinning, critical conditions for unpinning and vortex density distributions, with traps and vortex-free regions are in agreement with the framework of vortex creep theory.

The sudden increase in the crust rotation rate Ω_c and the decrease in the superfluid rotation rate Ω_s at a glitch offset the lag ω from its pre-glitch steady state value. If the creep process has a linear dependence on the lag, the response to the offset is simple exponential relaxation. Several components of exponential relaxation are observed in the Vela pulsar. After the exponential relaxation is over the relaxation of the glitch continues, actually until the time of the next glitch, in a characteristic non-exponential manner. The glitch in the spin-down rate recovers approximately linearly in time, i.e. to the lowest order, with a constant second derivative $\ddot{\Omega}$ of the rotation rate. Just as exponential relaxation is the signature of linear dynamics, power-law relaxation processes indicate nonlinear dynamics. It is an appealing feature of the vortex creep model that both types of observed post-glitch behaviour can occur, in different parts of the neutron star superfluid, as two regimes of the same physical process. Being a process of thermal activation, the creep process has an intrinsic exponential dependence on pinning energies, temperature and the driving lag ω through Boltzmann factors. Depending on the pinning energy E_p , the temperature and the steady spin-down rate $\dot{\Omega}_\infty$ dictated by the pulsar torque, the creep current in some parts of the superfluid can have a linear dependence on the lag, leading to exponential post-glitch response; while other parts of the superfluid have the full nonlinear dependence on the glitch induced perturbation (Alpar et al. 1989). In the nonlinear regime the steady state lag is very close to the critical lag, $\omega_{\text{cr}} - \omega_\infty \ll \omega_{\text{cr}}$. This makes it possible to reach the critical conditions for unpinning by fluctuations from steady state as supported by simulations (Melatos & Warszawski 2009).

The focus of this paper is on the nonlinear creep response by which we model the interglitch recovery at constant $\ddot{\Omega}$, after the exponential relaxation is over. When the superfluid rotation rate is reduced by $\delta\Omega$ as vortices unpin at the glitch move through a nonlinear creep region of moment of inertia δI , the nonlinear creep current, with its very sensitive dependence on the lag, will stop. This region is not spinning down after the glitch. As the pulsar torque is acting on less moment of inertia, the spin-down rate will increase




Figure 1. (a) Predicted ‘Fermi function behaviour’. Part of the glitch in the spin-down rate, $\Delta\dot{\Omega} = (\delta I/I)\dot{\Omega}$, recovers at time $t_0 = \delta\Omega/|\dot{\Omega}|$. This figure is taken from Alpar et al. (1984a). (b) The observed step recovery, at $t_0 \cong 400$ d after the 1988 glitch, is shown, superposed on the ‘triangle recovery’. This figure is taken from Buchner & Flanagan (2008).

by a step $\Delta\dot{\Omega}$, such that $\Delta\dot{\Omega}/\dot{\Omega} = \delta I/I$ where I is the total moment of inertia of the star. The lag will return to its pre-glitch value after a waiting time $t_0 \equiv \delta\Omega/|\dot{\Omega}|$, since the star continues to spin-down under the pulsar torque. Around time t_0 the creep will start again, showing up as a positive step recovery $|\Delta\dot{\Omega}| = |\dot{\Omega}|\delta I/I$, which will be observable if δI is large enough. This extreme nonlinear signature of stop-hold-and-restart was a prediction of the vortex creep model, dubbed ‘Fermi function behaviour’ (Alpar et al. 1984a). This has been observed clearly in one instance in the Vela pulsar (Flanagan 1995; Buchner & Flanagan 2008). Fig. 1 shows a sketch of the predicted behaviour and the observed step recovery for 400 d after the 1988 glitch of the Vela pulsar providing strong direct evidence for the presence of nonlinear creep.

More likely, the vortices unpin at the glitch will cause the superfluid rotation rate to decrease by amounts $\delta\Omega_i$ as they move through many nonlinear creep regions i with moments of inertia δI_i . Note that δI_i does not refer to an actual increase or decrease in the moment of inertia; it denotes a *small* amount of moment of inertia belonging to the creep region i where the superfluid rotation rate has decreased at the glitch by the particular amount $\delta\Omega_i$. At the time of

the glitch creep stops in all of these regions, causing a total increase in the spin-down rate by

$$\frac{\Delta\dot{\Omega}(0)}{\dot{\Omega}} = \frac{\sum_i \delta I_i}{I} \equiv \frac{I_A}{I}, \quad (1)$$

where I_A is the total moment of inertia of nonlinear creep regions through which vortices moved suddenly at the glitch. Each component of the inertial moment, δI_i , will restart creep at its own waiting time $t_i \equiv \delta\Omega_i/|\dot{\Omega}|$; these are not discernible as step recovery in the spin-down rate, as individual δI_i are too small. Instead a stacked response of the nonlinear recovery of creep regions with moments of inertia δI_i at successive times t_i will be observed. In particular, in the lowest, mean field approximation, if a uniform average area density of vortices unpins, moves through nonlinear creep regions of total moment of inertia I_A , and repins, the cumulative response is a ‘triangle recovery’ of the glitch step $\Delta\dot{\Omega}$, with a constant second derivative of the rotation rate

$$\ddot{\Omega} = \frac{I_A}{I} \frac{\dot{\Omega}^2}{\delta\Omega}, \quad (2)$$

where $\delta\Omega$ is the maximum glitch induced decrease in the superfluid rotation rate, corresponding to the total number of vortices involved in the glitch. This ‘triangle recovery’, as a lowest order approximate fit to the data has been observed to extend till the next glitch in every interglitch interval of the Vela pulsar. The end of the triangle response signals the return of the average vortex density to pre-glitch conditions. Assuming that glitches start at critical conditions involving vortex density, the recovery time

$$t_g \equiv \frac{\delta\Omega}{|\dot{\Omega}|} \equiv \frac{|\Delta\dot{\Omega}|}{\dot{\Omega}} \quad (3)$$

is roughly when the star is ready for the next glitch. Thus t_g provides a first estimate of the time to the next glitch. The triangle recovery can be written as

$$\frac{\Delta\dot{\Omega}(t)}{\dot{\Omega}} = \frac{I_A}{I} \left(1 - \frac{t}{t_g}\right). \quad (4)$$

To the extent that the density of unpinned vortices is not uniform, the observed post-glitch $\dot{\Omega}(t)$ will deviate from a linear time dependence (constant $\ddot{\Omega}$). The t^2 and higher order polynomial terms in $\dot{\Omega}(t)$ are small compared to the lowest order, constant $\ddot{\Omega}$ term. In our model this reflects the dominance of the uniform term in the spatial distribution of the unpinned vortex density $\delta n(r) \equiv \delta n_0$, higher moments (spatial fluctuations) of $\delta n(r)$ being weaker than the mean δn_0 .

Equations (1) and (2) are complemented by the angular momentum balance at the glitch when the motion of unpinned vortices leads to a reduction in the superfluid rotation rate with maximum value $\delta\Omega$. The glitch induced decrease in the superfluid rotation rate, $\delta\Omega_i$, varies linearly with δI_i , between 0 and the maximum value $\delta\Omega$, throughout the constituent small regions with moments of inertia δI_i that collectively make up the creep regions with total moment of inertia I_A . The average is $\delta\Omega/2$. The total angular momentum transfer in the glitch from the nonlinear creep regions in the superfluid to the crust is therefore $(1/2)I_A\delta\Omega$. This angular momentum is transferred back to the superfluid during the triangle recovery. The vortices unpinned at the glitch also move through the vortex-free (capacitor) regions, of total moment of inertia I_B , interspersed with the nonlinear creep regions, yielding an additional angular momentum transfer $I_B\delta\Omega$, which is not returned to the superfluid. This irreversible vortex discharge at the glitch is analogous to capacitor discharges. It accounts for the part of the glitch in frequency that

does not relax back, as is seen in the observations. The total angular momentum lost by these components of the superfluid accounts for the observed spin-up $\Delta\Omega_c$ that remains after the exponential relaxations:

$$I_c\Delta\Omega_c = \left(\frac{1}{2}I_A + I_B\right)\delta\Omega. \quad (5)$$

Equations (1), (2) and (5) constitute the nonlinear vortex creep model relating the long-term interglitch behaviour to the glitch parameters. Using the observed values of the glitch in the rotation frequency, $\Delta\Omega_c$, and in the spin-down rate, $\Delta\dot{\Omega}$, after the exponential relaxations are subtracted, together with the observed interglitch $\ddot{\Omega}$, these three equations can be solved for the model parameters I_A , I_B and $\delta\Omega$, or, equivalently, for t_g , the estimated time to the next glitch. We shall call this our Model (1). Note that the interglitch recovery depicted in equation (4) is by itself a model independent phenomenological description of the observed timing behaviour. It can be used as a basis for braking index calculations that take into account the full interglitch behaviour extending till the next glitch. The vortex creep model provides the physical context for understanding the power-law (constant $\ddot{\Omega}$) interglitch timing which is a clear signal of nonlinear dynamical behaviour. Moreover, the natural relation between the glitch parameters provided by the vortex creep model has been found to have remarkable predictive power for the time to the next glitch for the majority of glitch intervals, while the estimates can be improved by extending the vortex creep model.

Shannon et al. (2016) modelled the interglitch timing with exponential recovery with a constant relaxation time $\tau = 1600$ d, and ascribed the residuals to a noise process with a power-law spectrum. They find that each interglitch interval and the total spin-down rate data set have similar power spectra and conclude that the noise process is stationary. Their calculated spin-down rate time series however does not contain the frequency jumps at glitches, an essential ingredient in the vortex creep model for the determination of the subsequent second derivative $\ddot{\Omega}$ of the rotation rate. The fact that $\dot{\Omega}$ correlates with the glitch parameters in accordance with equations (1), (2) and (5) therefore favours the vortex creep model versus a noise process that can reproduce the same behaviour. Employing the standard formula for the braking index, $n \equiv \Omega\ddot{\Omega}/\dot{\Omega}^2$ to the constant $\ddot{\Omega}$ due to the internal torques leads to an ‘anomalous’ interglitch braking index given by equations (1), (2) and (5);

$$n_{\text{ig}} \equiv \frac{\Omega\ddot{\Omega}_{\text{int}}}{\dot{\Omega}^2} = \left(\frac{1}{2} + \frac{I_A}{I_B}\right) \left(\frac{\Delta\dot{\Omega}}{\dot{\Omega}}\right)^2 \left(\frac{\Delta\Omega}{\dot{\Omega}}\right)^{-1}_{-3} \left(\frac{\Delta\Omega}{\dot{\Omega}}\right)^{-1}_{-6}. \quad (6)$$

The subscripts -3 and -6 mean the corresponding dimensionless ratios are normalized to 10^{-3} and 10^{-6} , respectively.

3 OBSERVATIONS AND MODEL FITTING

Since 1985, the Vela pulsar has been observed on most days using the Hartebeesthoek Radio Astronomy Observatory (HartRAO) telescope. Observations are made at either 1668 or 2273 MHz using the 26 m telescope. The South polar bearing failed on 2008 October 3 and was repaired on 2010 July 22. During this period the telescope could only be driven in declination. Observations were made of Vela as it transited. The MeerKAT precursor XDM (now HartRAO 15 m) was used for Vela observations from 2009 Feb 6 until the telescope was repaired. The Tempo2 glitch plugin (Edwards, Hobbs & Manchester 2006; Hobbs, Edwards & Manchester 2006) was used to calculate $\dot{\Omega}$ values.

Alpar et al. (1984b) initially analysed the post-glitch behaviour of the Vela pulsar for its first four glitches within the vortex creep model. This work used data with major uncertainties in the actual dates of the glitches. Alpar et al. (1993) and Chau et al. (1993) then examined the post-glitch recovery of the first nine glitches of the Vela pulsar. They used the vortex creep model in which the post-glitch relaxation is described by the equation:

$$\frac{\Delta\dot{\Omega}_c(t)}{|\dot{\Omega}|_\infty} = -\sum_{i=1}^3 \frac{I_i}{I} \frac{\Delta\Omega_c(0)}{|\dot{\Omega}|_\infty \tau_i} e^{-t/\tau_i} - \frac{I_A}{I} + \frac{I_A}{I t_g} t. \quad (7)$$

The first three terms express the short-term exponential relaxation response to a glitch. The remaining two terms are relevant for the long time-scale and describe the nonlinear response. Exponential relaxation components with time-scales of hours, days and months are commonly observed after Vela pulsar glitches. Theoretical interpretations of the exponential relaxation terms include superfluid mutual friction (e.g. Haskell & Antonopoulou 2014), vortex creep in the linear regime Alpar et al. (1989), vortex-lattice interactions Jones (1993) and Ekman pumping van Eysden & Melatos (2010). Alpar et al. (1993) found that $\tau_1 = 10$ h, $\tau_2 = 3.2$ d and $\tau_3 = 32$ d, described all Vela glitches. The exponential relaxations were followed by the long-term triangle recovery of the spin-down rate from which they extracted estimates of the time t_g to the next glitch.

We follow the fitting procedure of the earlier applications to analyse the long-term relaxation of the 1994 double glitch and the 1996, 2000, 2004, 2006, 2010 and 2013 glitches. For these later glitches we use data from the HartRAO. We use the Levenberg–Marquardt method to find the best-fitting values with MPFITFUN procedure (Markwardt 2009).² A quick look at the data shows that by 100 d after each glitch all exponential relaxation components are fully relaxed (Yu et al. 2013). We therefore use data starting from 100 d after each glitch for our long-term interglitch fits with the last two terms in equation (7) which describe the ‘triangle’ nonlinear creep response, equation (4). The fits are shown in Fig. 2.

The best-fitting constant $\dot{\Omega}$ values, inferred parameters I_A/I , I_B/I and t_g and observed times t_{obs} to the next glitch are tabulated for the current sample of 17 glitches in Table 1 which presents the results of Alpar et al. (1993) for the first eight glitches, of Chau et al. (1993) for the ninth glitch and our results for the last eight glitches. Errors quoted for $\dot{\Omega}$ are formal linear regression errors, which propagate to give the same percentage errors in n_{ig} . For the most recent, 2013, glitch we have an estimated time to the next glitch, $t_g = 1553$ d, which gives the expected glitch date as 2017 December 22. The moment of inertia fractions I_1/I , I_2/I and I_3/I associated with exponential relaxations with $\tau_1 = 10$ hr, $\tau_2 = 3.2$ d and $\tau_3 = 32$ d are also tabulated. These are not relevant to the long-term recovery discussed in this paper. They do, however, contribute to lower bounds on the moment of inertia of creeping superfluid, which in turn are relevant to possible constraints on the equation of state of neutron star matter (Datta & Alpar 1993; Link, Epstein & Lattimer 1999). I_1/I , I_2/I and I_3/I are included in Table 1.

As a measure of the estimates we shall use

$$\Delta t_i \equiv t_{g,i} - t_{\text{obs},i}. \quad (8)$$


For the first nine glitches analysed earlier, the mean $\overline{\Delta t}(9) = 200$ d and the standard deviation $\sigma(9) = 321$ d. Interestingly, for seven out of the first nine glitches the estimator t_g was longer than t_{obs} . For the current sample the mean fractional deviation between the estimated

and observed intervals between the glitches is now $\overline{\Delta t}(15) = 132$ d and the standard deviation $\sigma(15) = 256$ d for the 15 interglitch intervals. For 11 out of the 15 glitches with observed times to the next glitch, the estimated t_g is longer than t_{obs} . As the sample of glitches with an observed time to the next glitch has increased from 9 to 15, the mean accuracy of the estimate has decreased from $\overline{\Delta t}(9) = 200$ d to $\overline{\Delta t}(15) = 132$ d, with the standard deviation decreasing from $\sigma(9) = 321$ d to $\sigma(15) = 256$ d. This strongly supports the validity of the nonlinear vortex creep model, in particular, to the association of the interglitch $\dot{\Omega}$ with a recovery process the completion of which indicates the re-establishment conditions prone to a glitch. It is significant that the mean offset of our estimates t_g from the observed glitch intervals t_{obs} remains positive, even though the sample almost doubled. Of 15 events in the present sample, 11 have $|\Delta t_i| < \overline{\Delta t}(15) = 132$ d. For these 11 interglitch intervals, the mean $\overline{\Delta t}(11) = 14.5$ d and $\sigma(11) = 131$ d only, so the model prediction is quite successful. The four events that have $|\Delta t_i| > \overline{\Delta t}(15) = 132$ d all have, suggestively, t_g significantly longer than t_{obs} ; 344 d $< \Delta t_i < 712$ d, with mean $\overline{\Delta t}(4) = 497$ d and $\sigma(4) = 197$ d. These four glitches will be addressed with an extension of the model, positing that these glitches incorporated ‘persistent shifts’ in the spin-down rate as commonly observed in the Crab pulsar’s glitches. If there is a persistent shift which is not taken into account, the estimated t_g would be systematically longer than the observed intervals t_{obs} , as detailed below, suggesting that the persistent shift may provide the explanation for why all four instances of $>1\sigma$ deviations have $t_g > t_{\text{obs}}$. Without a systematic reason, the random occurrence of $t_g > t_{\text{obs}}$ in these four instances has a probability of 1/16. The idea that this behaviour, common in the Crab pulsar glitches, can also occur in some occasional Vela glitches is consistent with the evolution of glitch behaviour with age as one progresses from the Crab pulsar to the Vela pulsar and older pulsars. In the next Section we shall describe and apply a new extended model for predicting the time to the next glitch.


4 THE MODIFIED INTERGLITCH TIMES OF THE VELA PULSAR

The fact that the four glitches whose observed arrival times t_{obs} show substantial discrepancies arrive earlier than the estimated times suggest that recovery is complete when the spin-down rate returns to a value somewhat less than its pre-glitch value. This suggests that each Vela pulsar glitch may be accompanied by a ‘persistent shift’, a part of the step in the spin-down rate which never recovers, as observed in the Crab pulsar’s glitches (Lyne et al. 2015). For the Crab pulsar, assuming that the glitches are pure unpinning events with creep response leads to estimated t_g of the order of a few months in disagreement with the observed glitch times (Alpar et al. 1996). This is interpreted as evidence that the glitches in the Crab pulsar are not due to vortex unpinning alone. Alpar et al. (1996) proposed that the comparatively small ($\Delta\Omega/\Omega \sim 10^{-8}$) and infrequent (~ 6 yr interglitch time intervals) events in Crab are starquakes (Baym & Pines 1971), in conjunction with unpinning events. In the framework of the vortex creep model, crust cracking is associated with stresses induced by pinned vortices. This scenario also explains the permanent offsets in the spin-down rate, the ‘persistent shifts’, observed in Crab pulsar glitches. The persistent change in spin-down rate is due to newly created vortex traps with surrounding vortex-free regions introduced by crust cracking. The newly created vortex-free regions were sustaining vortex creep and contributing to spin-down


² <http://purl.com/net/mpfit>




(a) 1994




(b) 1996




(c) 2000




(d) 2004



(e) 2006



(f) 2010



(g) 2013

Figure 2. Model fits to post-glitch spin-down rate after the 1994, 1996, 2000, 2004, 2006, 2010 and 2013 glitches. For the 1994 double glitch, the model fit is applied to data after the second event. The bottom panels show the residuals.

Table 1. The inferred and observed parameters for the Vela glitches. The entries for the first eight glitches and the ninth glitch are taken from Alpar et al. (1993) and Chau et al. (1993), respectively. Errors are given in parentheses. The last two columns give the interglitch constant $\dot{\Omega}$ and the anomalous braking index n_{ig} .

Yr	t_{obs} (d)	t_g (d)	Δt_i (d)	$(I_A/I)_{-3}$	$(I_B/I)_{-3}$	$(I_1/I)_{-3}$	$(I_2/I)_{-3}$	$(I_3/I)_{-3}$	$(\dot{\Omega})_{-21}(\text{rads}^{-3})$	n_{ig}
1969	912	1624	712	7.1	8.4	0.59	1.5	5.8	5.0	36.4
1971	1491	1375	-116	7.2	8.8	0.59	1.5	6.4	5.9	43.6
1975	1009	1036	27	7.2	12.4	0.59	1.5	5.1	7.9	57.8
1978	1227	1371	144	6.6	15.2	0.59	1.5	10.0	5.5	40.0
1981	272	616	344	6.3	12.2	0.59	1.5	3.2	1.16	85.0
1982	1067	1485	418	6.0	8.4	0.59	1.5	12.1	4.6	33.6
1985	1261	972	-289	6.5	7.9	0.59	1.5	9.0	7.6	55.6
1988	907	1422	515	4.7	8.2	0.59	1.5	9.5	3.7	27.5
1991	1102	1151	49	7.4	16.2	0.59	1.5	10.7	7.3	53.5
1994	778	765	-13	5.9(0.03)	18.8 (0.07)	0.59	1.5	8.3	8.7 (0.22)	64.1
1996	1190	1072	-118	6.6 (0.02)	13.1 (0.14)	0.59	1.5	12.4	7.0 (0.33)	51.2
2000	1634	1644	10	6.4 (0.02)	12.4 (0.03)	0.59	1.5	4.8	4.4 (0.21)	32.4
2004	767	913	146	6.7 (0.02)	15.8 (0.08)	0.59	1.5	7.6	8.3 (0.19)	61.0
2006	1449	1464	15	5.0 (0.01)	12.4 (0.11)	0.59	1.5	11.8	3.9 (0.23)	23.4
2010	1147	1281	134	6.1 (0.02)	25.2 (0.10)	0.59	1.5	9.2	5.4 (0.17)	39.6
2013	-	1553	-	6.4 (0.01)	13.4 (0.07)	0.59	1.5	6.9	4.7 (0.09)	34.2




Figure 3. The schematic view of the long-term behaviour of $\Delta\dot{\Omega}$.

before the glitch, while they no longer contribute to $\dot{\Omega}$ after it. The resulting permanent shift in the spin-down rate is given by

$$\frac{\Delta\dot{\Omega}_p}{\dot{\Omega}} = \frac{I_b}{I}, \quad (9)$$

where $\Delta\dot{\Omega}_p$ is the observed permanent change in $\dot{\Omega}$ and I_b is the moment of inertia of the newly created capacitor region. Crust cracking irreversibly restructures the vortex pinning distribution by introducing new capacitor elements. The steady state $\dot{\Omega}$ value for subsequent glitches is permanently reset to a new value that is less by $\Delta\dot{\Omega}_p$. The next glitch would then occur roughly when $\dot{\Omega}$ has returned to $\Delta\dot{\Omega}_p$ less than its steady state value $\dot{\Omega}_{n,-}$ before the previous glitch:

$$\dot{\Omega}_{n+1,-} = \dot{\Omega}_{n,-} - \Delta\dot{\Omega}_p. \quad (10)$$

If such persistent shifts also occur in the Vela pulsar, they would be unresolved in the observed total $\Delta\dot{\Omega}$ in a glitch. However, the post-glitch nonlinear recovery at constant $\dot{\Omega}$ would be completed earlier, as shown in Fig. 3. We now introduce a modified interglitch time estimator t'_g , with the hypothesis that some Vela glitches have a small ‘persistent shift’ as observed in the Crab pulsar.

In the standard vortex creep model (Alpar et al. 1996) in young pulsars like Crab, there is a building phase in which the vortex trap and the vortex-free regions surrounding them are being formed in each glitch, while this building phase is mostly over in the Vela pulsar. In the Crab and Vela pulsars glitches approximately the same number of vortices participate in the glitches. This number, the




Figure 4. Spin-down rates at the fiducial epochs t_{obs} (black points, Model 1), t'_g (red points, Model 2) and t'_g (green points, Model 3) prior to the next glitch (1971–2013). The best straight line fits used to extract the braking index n are also shown.

typical number of vortices unpinning collectively, then acts rather in the manner of a relay race. In the azimuthal direction, the unpinned vortices move rapidly relative to the crust lattice through vortex-free regions that surround the distribution of vortex traps, while scattering some small distance in the radially outward direction. By the time one batch of unpinned vortices have equilibrated with the superfluid flow they will have ‘passed the baton’ by causing the unpinning of a similar number of vortices in vortex traps that were close to critical conditions for unpinning. In the Crab (and other young pulsars) the vortex traps are yet sparse. A batch of unpinned vortices will affect a few other traps on their path and trigger these traps to unpin, but the avalanche will not travel far, as it fails to find more traps to unpin. So the total radial distance (or moment of inertia) that the relays of unpinned vortices travel through is only a small fraction of the moment of inertia in the crust superfluid. The size of the glitch depends on the total angular momentum transfer from the superfluid to the normal matter crust. Hence, the Crab glitches are much smaller than those in the Vela pulsar. The fact that older pulsars all exhibit glitches that are comparable in size

Table 2. Estimations of the model parameters for the four glitches for which t_g is considerably longer than t_{obs} if the persistent shift is the same magnitude as the average value for Crab, $\Delta\dot{\Omega}_{\text{per}}/\dot{\Omega} = 2.5 \times 10^{-4}$.

Yr	t_{obs} (d)	t'_g (d)	Δt_i (d)	$(I_A/I)_{-3}$	$(I_B/I)_{-3}$
1969	912	1567	655	6.9	9.0
1981	272	592	320	6.1	12.9
1982	1067	1423	356	5.8	9.1
1988	907	1346	439	4.5	8.9

to those of the Vela pulsar suggests that these glitches involve the entire crust superfluid, which represents a connected network for the unpinning vortices to percolate through in relays. The Crab pulsar glitches only involve a small segment of the crust lattice connected to the current site of vortex unpinning, so that the Crab pulsar glitches are not much larger than the minimum glitch size recently resolved (Espinoza et al. 2014). Thus we arrive at an evolutionary picture, with vortex traps formed in each successive glitch building up from disjoint segment networks of neighbouring vortex traps where unpinning events can travel, as in the Crab pulsar, into a connected network allowing unpinning events to percolate through the entire crust superfluid, defining the maximum glitch size, as observed in Vela and older pulsars. This is supported by the fact that Vela and older pulsars have occasional small Crab-sized glitches while all of the Crab pulsar glitches observed so far have magnitudes $\Delta\Omega/\Omega \leq 10^{-8}$. Having exhibited two distinct small, Crab-sized glitches, the Vela pulsar might also have Crab-like persistent shifts in the spin-down rate accompanying some of its glitches. Thus, we are now allowing for the possibility that the Vela pulsar may still be rearranging its network of vortex traps by forming a new vortex trap and surrounding vortex-free region during some glitches.

The part of the $\Delta\dot{\Omega}$ associated with the permanent shift cannot be discerned at the time of the glitch. Being a persistent shift, it will not relax back. The part of the step down in $\dot{\Omega}$ that restores with a constant $\dot{\Omega}$ will continue the recovery until new steady state conditions that include the persistent shift are reached. A schematic view of the long-term behaviour of $\dot{\Omega}(t)$ is depicted in Fig. 3. In models with persistent shifts, the triangle has less depth, and the estimated time of the next glitch is a bit shorter. This will make the predicted glitch times closer to the observed values. The new estimated glitch time intervals become

$$t'_g = t_g - \frac{|\Delta\dot{\Omega}_p|}{\dot{\Omega}} = \frac{|\Delta\dot{\Omega}|}{\dot{\Omega}} - \frac{|\Delta\dot{\Omega}_p|}{\dot{\Omega}}, \quad (11)$$

where $\Delta\dot{\Omega}$ and $\Delta\dot{\Omega}_p$ are the total and the permanent step in the spin-down rate, respectively.

The values of $\Delta\dot{\Omega}_p/\dot{\Omega}$ are between $(0.2 - 6.6) \times 10^{-4}$ for Crab pulsar glitches (Lyne et al. 2015). It may be different in Vela due to evolutionary reasons. We focus on the four glitches, for which the estimated time t_g to the next glitch was considerably longer than the observed interval t_{obs} . As Model (2), we assume that each of these

four glitches had a persistent shift of the same fractional magnitude as the average persistent shift for the Crab pulsar glitches.

$$\frac{\Delta\dot{\Omega}_p}{\dot{\Omega}} = \left\langle \frac{\Delta\dot{\Omega}_p}{\dot{\Omega}} \right\rangle_{\text{Crab}} = 2.5 \times 10^{-4}$$

The new estimates t'_g and Δt_i are given in Table 2. The mean and standard deviation for the four glitches are now $\overline{\Delta t}(4) = 443$ d and $\sigma(4) = 150$ d, while for the full sample of 15 glitches $\overline{\Delta t}(15) = 117$ d and the standard deviation $\sigma(15) = 241$ d. Introducing persistent shifts of the same magnitude as observed in the Crab pulsar is clearly not enough to make better estimates of the arrival times of the next glitch. We next consider the possibility that these five Vela glitches involved a major restructuring of the vortex trap network so that t_g was reduced by a third. The estimates $t'_g = (1 - 0.33)t_g$, Model (3), given in Table 3, lead to $\overline{\Delta t}(4) = 73$ d and $\sigma(4) = 152$ d, while for the full sample of 15 glitches $\overline{\Delta t}(15) = 18$ d and the standard deviation $\sigma(15) = 127$ d. Tables 2 and 3 also give the model parameters I_A/I and I_B/I . We have applied Kolmogorov–Smirnov (KS) tests to Model(1), without persistent shifts, and Model(3), with persistent shifts corresponding to a substantial rearrangement of the vortex creep regions. We first construct a cumulative distribution for the observed interglitch intervals t_{obs} , without binning. Comparing the cumulative distribution of our Model(1) estimates t_g with the t_{obs} cumulative distribution, we find that the null hypothesis is rejected at a confidence level of only 39 per cent. Comparing the cumulative distribution of the estimates t'_g of Model (3) with the cumulative t_{obs} distribution, a confidence level of 98 per cent is obtained. We conclude that our Model(3) with large persistent shifts produces substantial agreement with the observed glitch time intervals. The simulations of Melatos & Warszawski (2009) and Warszawski & Melatos (2011) also suggest that a substantial fraction of vortex traps and vortex-free regions are involved and rebuilt in each glitch.

5 THE BRAKING INDEX OF THE VELA PULSAR

We now use the above results to calculate the braking index for the Vela pulsar. Estimating the braking index requires modelling and subtracting all contributions of glitches and post-glitch and interglitch recovery. Lyne et al. (1996) did this by assuming that $\dot{\Omega}$ values 150 d after each glitch are already clean of post-glitch response and estimated a braking index $n \cong 1.4$. They used a post-glitch epoch when all short-term exponential relaxation components with $\tau_1 = 10$ h, $\tau_2 = 3.2$ d and $\tau_3 = 32$ d are over, but the interglitch recovery is far from finished. However, one also needs to take account of the constant $\dot{\Omega}$ response, which extends to the next glitch. The appropriate epochs when interglitch response is completed should be those that are just prior to the subsequent glitch. Espinoza et al. (2017) take into account the constant $\dot{\Omega}$ term, but assume that this recovery is not completed before the next glitch. The data fits in this work do not extend to the end of the interglitch interval. Instead fiducial epochs are chosen by extrapolating the $\dot{\nu}(t)$

Table 3. Estimations of the model parameters for the four glitches for which t_g is considerably longer than t_{obs} if $t'_g = (1 - 0.33)t_g$ (Model 3).

Yr	t_{obs} (d)	t'_g (d)	Δt_i (d)	$(\Delta\dot{\Omega}_{\text{per}}/\dot{\Omega})_{-3}$	$(I_A/I)_{-3}$	$(I_B/I)_{-3}$
1969	912	1088	176	2.3	4.8	15.5
1981	272	413	141	2.4	3.9	21.1
1982	1067	995	-72	2.3	3.7	15.3
1988	907	953	46	2.3	2.4	14.5

($\ddot{\Omega}$) fits back to immediately after the previous glitch. They obtain $n \cong 1.7 \pm 0.2$, and suggest that this low-braking index is due to incomplete recovery of the glitch in the spin-down rate. But the recovery continues up to, and is actually completed at, a time just before the next glitch.

In those glitches for which timing observations immediately before the next glitch are not available, we obtain the $\dot{\Omega}(t_{\text{obs},i})$ values at the epochs immediately prior to the next glitch by extrapolating from our fits with constant $\ddot{\Omega}$. We then produce the very long-term (between the years 1969 and 2013) $\ddot{\Omega}_{\text{PSR}}$ values for the Vela pulsar by the best linear fit to the $\dot{\Omega}(t_{\text{obs},i})$. This gives the estimate of $\ddot{\Omega}_{\text{PSR}}^{\text{obs}} = (3.83 \pm 0.15) \times 10^{-22} \text{ rad s}^{-3}$ leading to the braking PSR index $n_{t_{\text{obs}}} = 2.81 \pm 0.12$ for the Vela pulsar, with

$$n = \frac{\overline{\dot{\Omega}} \ddot{\Omega}_{\text{PSR}}}{(\overline{\dot{\Omega}})^2}, \quad (12)$$

where $\overline{\dot{\Omega}} = 70.4 \text{ rad s}^{-1}$ and $\overline{\ddot{\Omega}} = 9.8 \times 10^{-11} \text{ rad s}^{-2}$ are the average values over the 47 yr data span. The quoted errors in the braking index are propagated from the errors in our long-term fit for $\ddot{\Omega}_{\text{PSR}}$. The braking index $n_{t_{\text{obs}}} = 2.81 \pm 0.12$, using spin-down rates just prior to the observed time of the next glitch, is independent of any estimations of the glitch time. It is based only on the premises that the approximately constant second derivative of the rotation rate observed at interglitch epochs (i) is due to internal torques and (ii) determines the value of the spin-down rate just prior to the next glitch. This is supported by all the instances where timing data extend up to the next glitch. These data sets are well described by a constant $\ddot{\Omega}$ all the way to the next glitch, without any precursor signatures prior to the glitch. This justifies using extrapolations with the constant $\ddot{\Omega}$ in those interglitch data sets that do not extend to the next glitch.

Our other estimates of the braking index pick an estimated time of the next glitch (16 estimates including the next glitch after 2013) and are thus model dependent. Model (3) which gives better estimations of glitch times leads to $n_{t_g} = 2.87 \pm 0.17$. Within error bars, this is consistent with $n_{t_{\text{obs}}} = 2.81 \pm 0.12$.

6 CONCLUSIONS

We have confirmed that the interglitch timing behaviour of the Vela pulsar in the cumulative data from discovery to 2016, covering 16 glitches is characterized by a recovery of the spin-down rate at a constant $\ddot{\Omega}$, as was seen in the earlier data up to the ninth glitch. Power-law behaviour, like the constant $\ddot{\Omega}$, is a clear signal of nonlinear dynamics. This behaviour results from the cumulative manifestation of the step like ('Fermi function') recovery of the spin-down rate predicted by the vortex creep model (Alpar et al. 1984a) that has been directly observed in the Vela pulsar (Flanagan 1995; Buchner & Flanagan 2008). It is expected to determine the interglitch timing once the linear response in the form of exponential relaxations is over. The interglitch second derivative $\ddot{\Omega}$ of the rotation rate is determined by the parameters of the previous glitch. The completion of this ubiquitous interglitch behaviour signals the arrival of the next glitch and allows for an estimate of the time of the next glitch. We have shown that the constant behaviour dominates the interglitch interval for the full Vela pulsar data set comprising 16 glitches so far. Beyond the phenomenology, for 11 of the 15 intervals between glitches $\ddot{\Omega}$ relates to the glitch parameters in the way predicted by vortex creep theory, leading to predictions of interglitch intervals that agree with the observed glitch dates within

12 per cent. This level of agreement between theory and observation represents strong evidence in support of the nonlinear vortex creep theory.

The rms residuals from the constant $\ddot{\Omega}$ fits are $\lesssim 5$ per cent of $\Delta\dot{\Omega}$. These deviations correspond to spatial fluctuations from a uniform density of vortices unpinned in the glitch which gives rise to the constant $\ddot{\Omega}$ recovery.

With the daily monitoring of the Vela pulsar at HartRAO timing data extending to the next glitch are available for interglitch intervals since 1985, confirming the constant $\ddot{\Omega}$ behaviour. In those cases where there is a gap in the data prior to the next glitch, we have extrapolated the interglitch data to the epoch prior to the next glitch. Using observed or extrapolated values of the spin-down rate at the epochs prior to the glitches, we arrived at a braking index $n = 2.81 \pm 0.12$. This value is in agreement with the braking indices $n \leq 3$ observed for most isolated pulsars (Melatos 1997; Antonopoulou et al. 2015; Archibald et al. 2015; Lyne et al. 2015; Clark et al. 2016).

Furthermore, we find there is an explanation of the four cases for which the interglitch intervals predicted by the standard vortex creep theory are considerably longer than the observed intervals, with a mean fractional offset $\overline{\Delta t}(4) = 497 \text{ d}$ and standard deviation $\sigma(4) = 197 \text{ d}$. We propose that for each of these glitches there might be a persistent shift in spin-down rate that does not relax back entirely. This step which is commonly observed in the Crab pulsar glitches, would not be distinguished observationally as a part of the glitch in the spin-down rate. We introduce the possibility that some Vela glitches may involve the making of new vortex traps and vortex-free regions, as postulated earlier to explain the Crab pulsar's persistent shifts within vortex creep theory. After each glitch these newly restructured regions no longer contribute to spin-down and cause a permanent shift in the spin-down rate. The triangle interglitch behaviour in spin-down, which restores with a constant $\ddot{\Omega}$, is then completed somewhat earlier, restoring conditions for a new glitch. Deriving the modified estimator for the time between glitches by using this consideration, with persistent shifts of the average magnitude observed in the Crab pulsar, our Model (2), does not result in appreciable improvement of the estimates of the interglitch intervals. If we assume, as in Model (3), that as much as a third of the Vela pulsar's vortex creep network is restructured in each of these four glitches, the new interglitch time estimates after the four glitches have a mean fractional offset of only $\overline{\Delta t}(4) = 73 \text{ d}$ and standard deviation $\sigma(4) = 152 \text{ d}$ for the full sample of 15 glitches with an observed interval to the next glitch. It is suggestive that for all four glitches with t_g significantly longer than t_{obs} the idea that some glitches involve restructuring of the vortex creep network leads to better estimation. We do not have an understanding of why these particular four glitches involve rather substantial rearrangements, but this may be only natural in a complex system with nonlinear dynamics where the glitches are instabilities of the pinned vortex distribution, possibly triggered in interaction with lattice stresses. We use Models (2) and (3) as estimates of the range in predicted arrival times of the glitches, leading to a corresponding range of the braking index, $n = 2.93 \pm 0.15$ for Model (2) and $n = 2.87 \pm 0.17$ for Model (3). These are consistent, within error bars, with the value $n = 2.81 \pm 0.12$ obtained by using extrapolations to the epochs just prior to the next glitch.

We can now present predictions for the time of the glitch according to the vortex creep model. The next glitch will occur around $t_g = 1553 \pm 132 \text{ d}$ after the last observed glitch of 2013, around 2017 December 22, if no persistent shift was associated with the 2013 glitch. If a persistent shift of the order of those observed in

the Crab pulsar occurred then the glitch interval is $t_g = 1514 \pm 183$ d, predicting the arrival of the next glitch around 2017 November 11. If the 2013 glitch induced a large persistent shift associated with major restructuring of the vortex creep network, the next glitch will take place as early as 2016 July 27, within an interval of $t_g = 1040 \pm 152$ d. The uncertainties for each case correspond to one standard deviation.

An approximately constant second derivative of the rotation rate correlated with glitch parameters is also seen in the post-glitch or interglitch timing of most older pulsars which exhibit Vela-like giant glitches (Yu et al. 2013). Furthermore, the ‘anomalous’ braking indices observed from many pulsars (Johnston & Galloway 1999) can be understood as due to nonlinear creep response, with rotation rate second derivatives correlated with glitch parameters similar to those of the Vela pulsar (Alpar & Baykal 2006). The long-term interglitch timing of the Crab pulsar covering all glitches so far (Lyne et al. 2015; Čadež et al. 2016) and the large glitch associated change in the reported braking index of PSR J1846-0258 (Archibald et al. 2015) are also associated with the interglitch $\ddot{\Omega}$ predicted by the nonlinear vortex creep model, as we will discuss in future work.

Literally while we were submitting the first version of this paper we saw ATel # 9847 from Jim Palfreyman of the University of Tasmania, announcing that the Vela pulsar had glitched on that day, 2016 December 12 (Palfreyman 2016). The glitch has arrived 138 d late, within the 1σ uncertainty of 152 d from the prediction of our model with a persistent shift.

ACKNOWLEDGEMENTS

This work is supported by the Scientific and Technological Research Council of Turkey (TÜBİTAK) under the grant 113F354. MAA and DP thank the Aspen Center of Physics, which is supported by National Science Foundation grant PHY-1066293, for its hospitality during the period they collaborated on this paper, and MAA thanks the Simons Foundation for a grant that made his visit to the Aspen Center for Physics possible. MAA is a member of the Science Academy (Bilim Akademisi), Turkey. The data presented herein were obtained at the Hartebeesthoek Radio Astronomy Observatory (HartRAO). HartRAO is a National Facility of the National Research Foundation (NRF) of South Africa. We would like to thank the directors and staff of HartRAO for supporting the Vela monitoring program for the last 32 yr, and especially Claire Flanagan who started the Vela observing program and detected the first evidence for nonlinear response in the form of a step recovery of the spin-down rate. We thank Erbil Gügercinoğlu for useful discussions and Jim Palfreyman for alerting us to a mistake in the first version of the paper and for communications on the 2016 December 12 glitch.

REFERENCES

Alpar M. A., 1977, *ApJ*, 213, 527
 Alpar M. A., Baykal A., 1994, *MNRAS*, 269, 849
 Alpar M. A., Baykal A., 2006, *MNRAS*, 372, 489
 Alpar M. A., Anderson P. W., Pines D., Shaham J., 1984a, *ApJ*, 276, 325
 Alpar M. A., Anderson P. W., Pines D., Shaham J., 1984b, *ApJ*, 278, 791
 Alpar M. A., Cheng K. S., Pines D., 1989, *ApJ*, 346, 823
 Alpar M. A., Chau H. F., Cheng K. S., Pines D., 1993, *ApJ*, 409, 345
 Alpar M. A., Chau H. F., Cheng K. S., Pines D., 1996, *ApJ*, 459, 706

Anderson P. W., Itoh N., 1975, *Nature*, 256, 25
 Antonopoulou D., Weltevrede P., Espinoza C. M., Watts A. L., Johnston S., Shannon R. M., Kerr M., 2015, *MNRAS*, 447, 3924
 Archibald R. F., Kaspi V. M., Beardmore A. P., Gehrels N., Kennea J. A., 2015, *APJ*, 810, 67
 Baym G., Pines D., 1971, *Ann. Phys.*, NY, 66, 816
 Baym G., Pethick C., Pines D., 1969, *Nature*, 224, 673
 Buchner S. J., 2010, *Astron. Telegram*, 2768
 Buchner S., 2013, *Astron. Telegram*, 5406
 Buchner S., Flanagan C., 2008, in Bassa C., Wang Z., Cumming A., Kaspi V. M., eds., *AIP Conf. Proc. Ser. Vol. 983, 40 Years of Pulsars: Millisecond Pulsars, Magnetars and More*. Am. Inst. Phys., New York, p. 145
 Buchner S., Flanagan C., 2011, in Burgay M., D’Amico N., Esposito P., Pellizzoni A., Possenti A., eds., *AIP Conf. Proc. Ser. Vol. 1357, Radio Pulsars: An Astrophysical Key to Unlock the Secrets of the Universe*. Am. Inst. Phys., New York, p. 113
 Chau H. F., McCulloch P. M., Nandkumar R., Pines D., 1993, *ApJ*, 413, L113
 Clark C. J. et al., 2016, *ApJ*, 832, L15
 Datta B., Alpar M. A., 1993, *A&A*, 275, 210
 Dodson R. G., McCulloch P. M., Lewis D. R., 2002, *ApJ*, 564, L85
 Dodson R., Buchner S., Reid B., Lewis D., Flanagan C., 2004, *IAU Circ.*, 8370
 Edwards R. T., Hobbs G. B., Manchester R. N., 2006, *MNRAS*, 372, 1549
 Espinoza C. M., Antonopoulou D., Stappers B. W., Watts A., Lyne A. G., 2014, *MNRAS*, 440, 2755
 Espinoza C. M., Lyne A. G., Stappers B. W., 2017, *MNRAS*, 466, 147
 Flanagan C., 1991, *IAU Circ.*, 5311
 Flanagan C. S., 1995, in Alpar M. A., Kiziloglu U., van Paradijs J., eds., *NATO Adv. Sci. Inst. Ser. C, Vol. 450, The Lives of the Neutron Stars*. Kluwer, Dordrecht, The Netherlands, Boston, MA, p. 181
 Flanagan C. S., Buchner S. J., 2006, *Cent. Bur. Electron. Telegrams*, 595
 Flanagan C., McCulloch P. M., 1994, *IAU Circ.*, 6038
 Gügercinoğlu E., Alpar M. A., 2014, *ApJ*, 788, L11
 Haskell B., Antonopoulou D., 2014, *MNRAS*, 438, L16
 Hobbs G. B., Edwards R. T., Manchester R. N., 2006, *MNRAS*, 369, 655
 Johnston S., Galloway D., 1999, *MNRAS*, 306, L50
 Jones P. B., 1993, *MNRAS*, 263, 619
 Link B., Epstein R. I., Lattimer J. M., 1999, *Phys. Rev. Lett.*, 83, 3362
 Lyne A. G., Pritchard R. S., Graham-Smith F., Camilo F., 1996, *Nature*, 381, 497
 Lyne A. G., Jordan C. A., Graham-Smith F., Espinoza C. M., Stappers B. W., Weltevrede P., 2015, *MNRAS*, 446, 857
 Markwardt C. B., 2009, in Bohlender D. A., Durand D., Dowler P., eds, *ASP Conf. Ser. Vol. 411, Astronomical Data Analysis Software and Systems XVIII*. Astron. Soc. Pac., San Francisco, p. 251
 McCulloch P. M., Klekociuk A. R., Hamilton P. A., Royle G. W. R., 1987, *Aust. J. Phys.*, 40, 725
 Melatos A., 1997, *MNRAS*, 288, 1049
 Melatos A., Warszawski L., 2009, *ApJ*, 700, 1524
 Mochizuki Y., Izuyama T., 1995, *ApJ*, 440, 263
 Packard R. E., 1972, *Phys. Rev. Lett.*, 28, 1080
 Palfreyman J., 2016, *Astron. Telegram*, 9847
 Radhakrishnan V., Manchester R. N., 1969, *Nature*, 222, 228
 Reichley P. E., Downs G. S., 1969, *Nature*, 222, 229
 van Eysden C. A., Melatos A., 2010, *MNRAS*, 409, 1253
 Wang N., Manchester R. N., Pace R. T., Bailes M., Kaspi V. M., Stappers B. W., Lyne A. G., 2000, *MNRAS*, 317, 843
 Warszawski L., Melatos A., 2011, *MNRAS*, 415, 1611
 Yu M. et al., 2013, *MNRAS*, 429, 688
 Čadež A., Zampieri L., Barbieri C., Calvani M., Naletto G., Barbieri M., Ponikvar D., 2016, *A&A*, 587, A99

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.